

Constructive Logic (15-317), Fall 2017

Assignment 0: Say hi to logic!

Contact: 15-317 Course Staff

Due Tuesday, September 5, 2017

Welcome to 15-317, Fall 2017 edition! In this introductory homework assignment, you will practice some basic principles you'll need for the rest of the course. As a special exception to the usual rules, for this homework assignment, you may collaborate with other students in the class on the answers to all of the questions as long as you do your write-up individually.

We STRONGLY SUGGEST that you typeset this homework assignment in L^AT_EX so that you learn how to typeset your proofs now, while the problems are easier and you are perhaps less busy. You can find the code handout for this assignment on autolab; you can use it as a starting point.

This assignment is due at the beginning of class on the above date and it must be submitted electronically on autolab. The link to the course's autolab page can be found on the course homepage once it becomes available. Submit your homework as a tar archive containing two files: `hw0.pdf` (your written solutions) and `hw0.tut` (your Tutch solutions).

One and one is one¹

Prove the following formulas using the inference rules given in lecture (be sure to name each rule when you use it).

Task 1 (1 point).

$$(A \wedge (A \supset B)) \supset B$$

Task 2 (2 points).

$$(A \wedge ((A \wedge A) \supset B)) \supset B$$

Task 3 (2 points).

$$(A \wedge (A \supset B)) \supset (B \wedge B)$$

¹For the appropriate definition of "and".

Doing constructive mathematics

An integer a is said to *divide* an integer b , written $a \mid b$, if there exists an integer k such that $b = ak$. We write $a \nmid b$ for $\neg(a \mid b)$.

Task 4 (1 point). Give an (informal) constructive proof of the following proposition: for all integers a , b , and c , if $a \mid b$ and $b \mid c$, then $a \mid c$.

Task 5 (1 point). Give an (informal) non-constructive proof of the following proposition: for all integers a , b , and c , one of the following is true:

- $a \nmid b$ or $b \nmid c$;
- $a \mid c$.

Hint: Use the proposition from the previous task.

Task 6 (1 point). In the previous task, you likely used the following instance of the law of the excluded middle

$$a \mid b \vee a \nmid b.$$

Though constructivism rejects the law of the excluded middle as a general axiom, it will still be the case that for some specific propositions A we have

$$(A \vee \neg A) \text{ true.}$$

In particular, this will be the case whenever A is decidable, that is, whenever there exists an effective procedure for deciding whether or not A is true. Give a brief and informal constructive justification for concluding $a \mid b$ when $\neg(a \nmid b)$.

The *Fundamental Theorem of Arithmetic* states that every integer greater than 1 either is prime or factors as the product of primes numbers, and moreover, that this factorisation is unique up to reordering of factors.

Task 7 (1 point). Is the following proof that $3 \nmid 10$ constructive? Justify your answer.

Proof. Assume to the contrary that $3 \mid 10$. Then there exists a k such that $10 = 3k$. By the fundamental theorem of arithmetic, k has some unique prime factorisation $k = \prod_{i=1}^n p_i$. So 10 factors into primes as $10 = 3 \prod_{i=1}^n p_i$. But we also know that 10 factors into primes as $10 = 2 \times 5$. The existence of two distinct prime factorisations for 10 contradicts the uniqueness guaranteed by the fundamental theorem of arithmetic. We thus conclude that $3 \nmid 10$. $\square$

Say hi to tutch!

In this homework you will be introduced to the proof checker *tutch*. If you were ever wondering whether using that inference rule was quite right or not, wonder no more! Tutch can check the correctness of your natural deduction proofs².

In order to use tutch, you have several options:

1. Use it on Andrew machines via the command:
`/afs/andrew/course/15/317/bin/tutch <file>.`
2. Use it on your own system if you have AFS configured, using the above command. Make sure your system's `CellAlias` file contains the entry `andrew.cmu.edu andrew` so that the symlink pointing from `/afs/andrew` to `/afs/andrew.cmu.edu` gets created. On Linux systems using OpenAFS, this file can usually be found at `/etc/openafs/CellAlias`.
3. Install your own local copy of tutch. See:
<http://www2.tcs.uni-lmu.de/~abel/tutch/>.

Task 8 (1 point). Type the following proof in a file named `hw0.tut`, check it with `tutch` and submit it with your written solutions on autolab:

```
proof andComm: A & B => B & A =
begin
  [ A & B;
    A;
    B;
    B & A ];
  A & B => B & A
end;
```

You can also run `tutch -r ./hw0.req hw0.tut` to compare your solution against the requirements file `hw0.req` provided with this assignment.

²Provided that you type them in the correct syntax.