Systems of Nonlinear Equations

Paul Heckbert

Computer Science Department
Carnegie Mellon University

Nonlinear Systems

One dimensional:

$f : \mathrm{R} \rightarrow \mathrm{R}$

e.g. $f(x) = ax-b = 0$ (linear)

or $f(x) = \sin(2x)-x = 0$ (nonlinear)

Multidimensional

$f : \mathrm{R}^n \rightarrow \mathrm{R}^n$

e.g. $f(x) = Ax-b = 0$ (linear system)

or $\{x^2+y^2+xy-1=0,\ x^2+y^2-xy-1=0\}$ (nonlinear: two ellipses)

also known as root finding

Four 1-D Root-Finding Methods

- Bisection Method
- Newton's Method
- Secant Method
- Linear Fractional Interpolation

Bisection Method (Binary Search)

Given $f()$, tol, and $[a,b]$ such that $\mathrm{sign}(f(a)) \neq \mathrm{sign}(f(b))$:

while $b-a>\mathrm{tol}$

$m = (a+b)/2$ midpoint

if $\mathrm{sign}(f(a)) = \mathrm{sign}(f(m))$: then

$a = m$ recurse on right half

else

$b = m$ recurse on left half

Guaranteed convergence! (if you can find initial a,b)

but only at a linear rate: $|\mathrm{error}_{k+1}| \leq c|\mathrm{error}_k|$, $c<1$

Newton's Method

Given $f()$, $f'()$, tol, and initial guess x_0
$k = 0$
do
  $x_{k+1} = x_k - f(x_k)/f'(x_k)$
  k++
while $x_k - x_{k-1} >$ tol

Can diverge (especially if $f' \approx 0$).
If it converges, does so at quadratic rate: $|e_{k+1}| \le c|e_k|^2$.
Requires derivative computation.

Secant Method

Like Newton, but approximates slope using point pairs

Given $f()$, tol, and initial guesses x_0, x_1
$k = 1$
do
  $x_{k+1} = x_k - f(x_k)(x_k - x_{k-1})/(f(x_k)-f(x_{k-1}))$
  k++
while $x_k - x_{k-1} <$ tol

Can diverge.
If it converges, does so at rate 1.618: $|e_{k+1}| \le c|e_k|^{1.618}$.

Linear Fractional Interpolation

Instead of linear approximation, fit with fractional linear approximation: $f(x) \approx (x-u)/(vx-w)$ for some u,v,w

Given $f()$, tol, and initial guesses a, b, c, $f_a=f(a)$, $f_b=f(b)$
do
  $h = (a-c)(b-c)(f_a-f_b)f_c / [(a-c)(f_c-f_b)f_a-(b-c)(f_c-f_a)f_b]$
  $a = b;\ f_a = f_b$
  $b = c;\ f_b = f_c$
  c += $h;\ f_c = f(c)$
while $h >$ tol

If it converges, does so at rate 1.839: $|e_{k+1}| \le c|e_k|^{1.839}$.

1-D Root-Finding Summary

- Bisection: safe: never diverges, but slow.
- Newton's method: risky but fast!
- Secant method & linear fractional interpolation : less risky, mid-speed.

- Hybrids: use a safe method where function poorly behaved, Newton's method where it's well-behaved.

Multidimensional Newton's Method

f and *x* are *n*-vectors

First order Taylor series approx.: $f(x+h) \approx f(x)+f'(x)h$

$$f = \begin{bmatrix} f_1 \\ \vdots \\ f_n \end{bmatrix} \quad x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \quad J_f(x) = f'(x) = \begin{bmatrix} \partial f_1/\partial x_1 & \cdots & \partial f_1/\partial x_n \\ \vdots & \ddots & \vdots \\ \partial f_n/\partial x_1 & \cdots & \partial f_n/\partial x_n \end{bmatrix}$$

$f(x+h)=0$ implies $h=-J^{-1}f$ (don't compute J^{-1}, but solve $Jh=f$)

so iteration should be

$$x_{k+1} = x_k - J^{-1}(x_k)f(x_k)$$

Method requires calculation of Jacobian – can be expensive

Example: Intersection of Ellipses

2 equations in 2 unknowns:

$$x^2+y^2+xy-1=0$$

$$x^2+y^2-xy-1=0$$