

# DEFORMING THE HIPPOCAMPAL MAP

David S. Touretzky<sup>1</sup>, Wendy E. Weisman<sup>2</sup>, Mark C. Fuhs<sup>1</sup>, William E. Skaggs<sup>3</sup>,  
Andre A. Fenton<sup>4,5</sup>, and Robert U. Muller<sup>5,6</sup>

<sup>1</sup> Computer Science Department & Center for the Neural Basis of Cognition, Carnegie Mellon University, Pittsburgh, PA

<sup>2</sup> 300 Woodlawn Avenue, Saint Paul, MN

<sup>3</sup> Arizona Research Laboratories, Division of Neural Systems, Memory, and Aging, Tucson, AZ

<sup>4</sup> Institute of Physiology, Academy of Sciences of the Czech Republic, Vídenská 1083, 142 20 Prague 4, Czech Republic

<sup>5</sup> Dept. of Physiology and Pharmacology, State University of New York Health Sciences Center at Brooklyn, Brooklyn, NY

<sup>6</sup> Medical Research Council Center for Synaptic Plasticity, Dept. of Anatomy, University of Bristol, Bristol UK

Number of words in Abstract: 182

Number of text pages: 35

Number of figures: 10

Number of tables: 2

Corresponding author:

David S. Touretzky  
Computer Science Department  
Carnegie Mellon University  
Pittsburgh, PA 15213-3981

phone: 412-268-7561  
fax: 412-268-3608  
e-mail: dst@cs.cmu.edu

## Funding Sources:

- National Institutes of Health award MH59932 to DST and WES
- National Science Foundation REU supplement to award IIS-9978403 to DST (to fund Wendy Weisman)
- National Science Foundation IGERT training grant DGE-9987588 to DST (support for Mark Fuhs)
- National Institutes of Health awards NS20686 and NS 37150 to RUM
- Medical Research Council (UK) Grant to RUM

## Abstract

To investigate conjoint stimulus control over place cells, Fenton et al. (2000a) recorded while rats foraged in a cylinder with  $45^\circ$  white and black cue cards on the wall. Card centers were  $135^\circ$  apart. In probe trials the cards were rotated together or apart by  $25^\circ$ . Firing field centers shifted during these trials, stretching and shrinking the cognitive map. Fenton et al. (2000b) described this deformation with an *ad hoc* vector field equation.

We consider what sorts of neural network mechanisms might be capable of accounting for their observations. In an abstract, maximum likelihood formulation, the rat's location is estimated by a conjoint probability density function of landmark positions. In an attractor neural network model, recurrent connections produce a bump of activity over a 2D array of cells; the bump's position is influenced by landmark features such as distances or bearings. If features are chosen with appropriate care, the attractor network and maximum likelihood models yield similar results, in accord with previous demonstrations that recurrent neural networks can efficiently implement maximum likelihood computations (Pouget et al., 1998; Deneve et al., 2001).

**Keywords:** place cell, attractor network, maximum likelihood, map deformation, rodent hippocampus