



Counting I: One To One Correspondence and Choice Trees



How many seats in this auditorium?



Hint:
Count without counting!



If I have 14 teeth on the top and 12 teeth on the bottom, how many teeth do I have in all?



Addition Rule

Let A and B be two disjoint finite sets.

The size of $A \cup B$ is the sum of the size of A and the size of B .


$$|A \cup B| = |A| + |B|$$

Corollary (by induction)

Let $A_1, A_2, A_3, \dots, A_n$ be disjoint, finite sets.

$$\left| \bigcup_{i=1}^n A_i \right| = \sum_{i=1}^n |A_i|$$

Suppose I roll a
white die and a black die.



$S \equiv$ Set of all outcomes where the dice show different values.

$$|S| = ?$$

$$36 - 6 = 30$$

↑
doubles

$$\underbrace{5 + 5 + \dots + 5}_{6 \text{ times}} = 30$$

$S \equiv$ Set of all outcomes where the dice show different values.

$$|S| = ?$$

$A_i \equiv$ set of outcomes where the black die says i and the white die says something else.

$$|S| = \left| \bigcup_{i=1}^6 A_i \right| = \sum_{i=1}^6 |A_i| = \sum_{i=1}^6 5 = 30$$

$S \equiv$ Set of all outcomes where the dice show different values.

$$|S| = ?$$

$T \equiv$ set of outcomes where dice agree.

$$|S \cup T| = \# \text{ of outcomes} = 36$$

$$|S| + |T| = 36 \quad |T| = 6$$

$$|S| = 36 - 6 = 30$$

$S \equiv$ Set of all outcomes where the black die shows a smaller number than the white die. $|S| = ?$

$A_i \equiv$ set of outcomes where the black die says i and the white die says something larger.

$$S = A_1 \cup A_2 \cup A_3 \cup A_4 \cup A_5 \cup A_6$$

$$|S| = 5 + 4 + 3 + 2 + 1 + 0 = 15$$

$S \equiv$ Set of all outcomes where the black die shows a smaller number than the white die. $|S| = ?$


$L \equiv$ set of all outcomes where the black die shows a **larger** number than the white die.

$$|S| + |L| = 30$$

It is clear by **symmetry** that $|S| = |L|$.


Therefore $|S| = 15$

It is clear by symmetry that $|S| = |L|$.



Pinning down the idea of symmetry by exhibiting a correspondence.

Let's put each outcome in S in correspondence with an outcome in L by *swapping* the color of the dice.



Pinning down the idea of symmetry by exhibiting a correspondence.

Let's put each outcome in S in correspondence with an outcome in L by **swapping** the color of the dice.

Each outcome in S gets matched with exactly one outcome in L , with none left over.

$$\text{Thus: } |S| = |L|.$$

Let $f:A \rightarrow B$

be a function from a set A to a set B .

f is **1-1** if and only if

$$\forall x, y \in A, x \neq y \Rightarrow f(x) \neq f(y)$$

f is **onto** if and only if

$$\forall z \in B \exists x \in A f(x) = z$$

Let $f:A \rightarrow B$

be a function from a set A to a set B .

f is **1-1** if and only if

$$\forall x, y \in A, x \neq y \Rightarrow f(x) \neq f(y)$$



f is **onto** if and only if

$$\forall z \in B \exists x \in A f(x) = z$$


There
Exists

For Every


Let's restrict our attention to
finite sets.




$$\exists 1-1 f:A \rightarrow B \Rightarrow |A| \leq |B|$$




\exists onto $f:A \rightarrow B \Rightarrow |A| \geq |B|$



\exists 1-1 onto $f:A \rightarrow B \Rightarrow |A| = |B|$



1-1 Onto Correspondence (just "correspondence" for short)




Correspondence Principle

If two finite sets can be placed into 1-1 onto correspondence, then they have the same size.

Correspondence Principle

If two finite sets can be placed into 1-1 onto correspondence, then they have the same size.



It's one of
the most
important
mathematical
ideas of
all time!

2 CS?


Question: How many n-bit sequences are there?

000000	β à	0
000001	β à	1 ←
000010	β à	2
000011	β à	3
	...	
1...11111	β à	2^{n-1}

2^n sequences

$S = \{a,b,c,d,e\}$ has many subsets.

$\{a\}, \{a,b\}, \{a,d,e\}, \{a,b,c,d,e\},$
 $\{e\}, \emptyset, \dots$



The empty set is a set with all the rights and privileges pertaining thereto.

Question: How many subsets can be formed from the elements of a 5-element set?

a	b	c	d	e
0	1	1	0	1

{ b c e }

1 means "TAKE IT"
0 means "LEAVE IT"

Question: How many subsets can be formed from the elements of a 5-element set?

a	b	c	d	e
0	1	1	0	1

Each subset corresponds to a 5-bit sequence (using the "take it or leave it" code)

$$S = \{a_1, a_2, a_3, \dots, a_n\}$$

$$b = b_1 b_2 b_3 \dots b_n$$

a_1	a_2	a_3	\dots	a_n
b_1	b_2	b_3	\dots	b_n

$$f(b) = \{a_i \mid b_i = 1\}$$

a_1	a_2	a_3	\dots	a_n
b_1	b_2	b_3	\dots	b_n

$$f(b) = \{a_i \mid b_i = 1\}$$


f is 1-1: Any two distinct binary sequences b and b' have a position i at which they differ. Hence, $f(b)$ is not equal to $f(b')$ because they disagree on element a_i .

a_1	a_2	a_3	\dots	a_n
b_1	b_2	b_3	\dots	b_n

$$f(b) = \{a_i \mid b_i = 1\}$$

f is onto: Let S be a subset of $\{a_1, \dots, a_n\}$. Let $b_k = 1$ if a_k in S ; $b_k = 0$ otherwise. $f(b_1 b_2 \dots b_n) = S$.

The number
of subsets of
an n -element
set is 2^n .



Let $f:A \rightarrow B$

be a function from a set A to a set B .

f is **1-1** if and only if

$$\forall x, y \in A, x \neq y \Rightarrow f(x) \neq f(y)$$

f is **onto** if and only if

$$\forall z \in B \exists x \in A f(x) = z$$

Let $f:A \rightarrow B$

be a function from a set A to a set B .

f is a 1 to 1 correspondence iff

$\forall z \in B \exists$ exactly one $x \in A$ s.t. $f(x)=z$

f is a k to 1 correspondence iff

$\forall z \in B \exists$ exactly k $x \in A$ s.t. $f(x)=z$



To count the number of horses in a barn, we count the number hoofs and then divide by 4.




If Finite set A
has a k to 1
correspondence
to finite set B,
then $\#B = \#A/k$





I own 3 beanies and 2 ties.
How many different ways can
I dress up in a beanie and a
tie?



A restaurant has a menu with
5 appetizers, 6 entrees, 3 salads,
and 7 desserts.

How many items on the menu?

- $5 + 6 + 3 + 7 = 21$

How many ways to choose a complete
meal?

- $5 * 6 * 3 * 7 = 630$

A restaurant has a menu with 5 appetizers, 6 entrees, 3 salads, and 7 desserts.

How many ways to order a meal if I might not have some of the courses?

- $6 * 7 * 4 * 8 = 1344$

Hobson's restaurant has only 1 appetizer, 1 entree, 1 salad, and 1 dessert.

2^4 ways to order a meal if I might not have some of the courses.

Same as number of subsets of the set
{Appetizer, Entrée, Salad, Dessert}


Leaf Counting Lemma

Let T be a depth n tree when each node at depth $0 \leq i \leq n-1$ has P_{i+1} children.

The number of leaves of T is given by:


$$P_1 P_2 \dots P_n$$

Choice Tree for 2^n n-bit sequences




We can use a "choice tree" to represent the construction of objects of the desired type.

2^n n-bit sequences



Label each leaf with the object constructed by the choices along the path to the leaf.



2 choices for first bit


X 2 choices for second bit

X 2 choices for third bit


...

X 2 choices for the n^{th}

Choice Tree




A choice tree is a rooted, directed tree with an object called a "choice" associated with each edge and a label on each leaf.



A choice tree provides a "choice tree representation" of a set S , if

- 1) Each leaf label is in S , and each element of S is some leaf label
- 2) No two leaf labels are the same



We will now combine
the correspondence
principle with the
leaf counting lemma
to make a powerful
counting rule for
**choice tree
representation.**

Product Rule

IF S has a choice tree representation with P_1 possibilities for the first choice, P_2 for the second, and so on,

THEN

there are $P_1 P_2 P_3 \dots P_n$ objects in S

Proof: The leaves of the choice tree are in 1-1 onto correspondence with the elements of S .

Product Rule

Suppose that all objects of a type S can be constructed by a sequence of choices with P_1 possibilities for the first choice, P_2 for the second, and so on.

IF

1) Each sequence of choices constructs an object of type S

AND

2) No two different sequences create the same object

THEN

there are $P_1 P_2 P_3 \dots P_n$ objects of type S .

How many different orderings of deck with 52 cards?

What type of object are we making?

- Ordering of a deck

Construct an ordering of a deck by a sequence of 52 choices:

52 possible choices for the first card;

51 possible choices for the second card;

50 possible choices for the third card;

...

1 possible choice for the 52^{cond} card.

How many different orderings of deck with 52 cards?

By the product rule:

$$52 * 51 * 50 * \dots * 3 * 2 * 1 = 52!$$


52 "factorial" orderings

Stirling:

$$\log n! \sim n \log n - n$$
$$\log 52! \sim 52 \log 52 - 52 \approx 153$$


A permutation or arrangement of n objects is an ordering of the objects.

The number of permutations of n distinct objects is $n!$



How many sequences of 7
letters are there?


$$26^7$$



How many sequences of 7 letters contain at least two of the same letter?

$$26^7 - 26 * 25 * 24 * 23 * 22 * 21 * 20$$

Sometimes it is easiest to count the number of objects with property Q , by counting the number of objects that do not have property Q .



A formalization

Let $S(x): \Sigma^* \rightarrow \{\text{True}, \text{False}\}$ be any predicate.

We can associate S with the set:

$$\text{OBJECTS}_S = \{x \in \Sigma^* \mid S(x)\}$$

the "object space" S (or objects of type S)

When OBJECTS_S is finite, let us define

$$\#\text{OBJECTS}_S = \text{the size of } \text{OBJECTS}_S$$

In fact, define $\#S$ as $\#\text{OBJECTS}_S$

Object property Q on object space S

Consider $Q(x): \text{OBJECTS}_S \rightarrow \{\text{True}, \text{False}\}$


Define $\neg Q(x): \text{OBJECTS}_S \rightarrow \{\text{True}, \text{False}\}$
As Input(x); return NOT Q(x)

Proposition: $\#Q = \#S - \#(\neg Q)$

How many of our objects
have property Q in object
space S ?

$\#Q$

$$= \#\text{OBJECTS}_S - \#(\neg Q)$$



Helpful Advice:

In logic, it can be useful to represent a statement in the contrapositive.

In counting, it can be useful to represent a set in terms of its complement.



If 10 horses race, how many orderings of the top three finishers are there?


$$10 * 9 * 8 = 720$$

The number of ways of ordering, permuting, or arranging r out of n objects.

n choices for first place, $n-1$ choices for second place, ...

$$n * (n-1) * (n-2) * \dots * (n-(r-1))$$

$$= \frac{n!}{(n-r)!}$$



Ordered Versus Unordered

From a deck of 52 cards how many **ordered** pairs can be formed?

- $52 * 51$

How many **unordered** pairs?

- $52*51 / 2$ ß divide by overcount

Each unordered pair is listed twice on a list of the ordered pairs, but we consider the ordered pairs to be the same.

Ordered Versus Unordered

From a deck of 52 cards how many **ordered** pairs can be formed?

- $52 * 51$

How many **unordered** pairs?

- $52*51 / 2$ ß divide by overcount

We have a 2 to 1 map from ordered pairs to unordered pairs. Hence: the
 $\#$ unordered pairs = ($\#$ ordered pairs)/2

Ordered Versus Unordered

From a deck of 52 cards how many **ordered** 5 card sequences can be formed?

- $52 * 51 * 50 * 49 * 48$

How many orderings of 5 cards?

- $5!$

How many **unordered** 5 card hands?

- $52 * 51 * 50 * 49 * 48 / 5! = 2,598,960$

A combination or choice of r out of n objects is an (unordered) set of r of the n objects.

The number of r combinations of n objects:


$$\frac{n!}{r!(n-r)!} = \binom{n}{r}$$

n choose r



The number of subsets of size r that can be formed from an n -element set is:

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}$$



How many 8 bit sequences
have 2 0's and 6 1's?

Tempting, but incorrect:

8 ways to place first 0, times

7 ways to place second 0

Violates condition 2 of product rule!

Choosing position i for the first 0 and
then position j for the second 0 gives
the same sequence as choosing position
 j for the first 0 and position i for the
second.

How many 8 bit sequences
have 2 0's and 6 1's?

1) Choose the set of 2 positions to put
the 0's. The 1's are forced.

$$\binom{8}{2} \times 1 = \binom{8}{2}$$

2) Choose the set of 6 positions to put
the 1's. The 0's are forced.

$$\binom{8}{6} \times 1 = \binom{8}{6}$$

Symmetry in the formula:

$$\binom{n}{r} = \frac{n!}{r!(n-r)!} = \binom{n}{n-r}$$

How many hands have at least 3
aces?

$\binom{4}{3} = 4$ ways of picking 3 of the 4 aces.

$\binom{49}{2} = 1176$ ways of picking 2 cards from the remaining 49 cards.

$$4 \times 1176 = 4704$$

How many hands have at least 3 aces?

How many hands have exactly 3 aces?

$$\binom{4}{3} = 4 \text{ ways of picking 3 of the 4 aces.}$$

$$\binom{48}{2} = 1128 \text{ ways of picking 2 cards non-ace cards.}$$

$$4 \times 1128 = 4512$$

How many hands have exactly 4 aces?


$$\binom{4}{4} = 1 \text{ way of picking 4 of the 4 aces.}$$

48 ways of picking one of the remaining cards

$$4512 + 48 = 4560$$

4704 \neq 4560

At least one of the
two counting
arguments is not
correct.



Four different sequences of choices produce the same hand

$\binom{4}{3} = 4$ ways of picking 3 of the 4 aces.

$\binom{49}{2} = 1176$ ways of picking 2 cards from the remaining 49 cards.

$$4 \times 1176 = 4704$$

A♣ A♦ A♥

A♠ K♦

A♣ A♦ A♠

A♥ K♦

A♣ A♠ A♥

A♦ K♦

A♠ A♦ A♥

A♣ K♦

Is the other
argument
correct? How do
I avoid fallacious
reasoning?



The Sleuth's Criterion

Condition (2) of the product rule:

For any object it should be possible to reconstruct the sequence of choices which lead to it.

1) Choose 3 of 4 aces

2) Choose 2 of the remaining cards

A♣ A♦ A♥ A♠ K♦

Sleuth can't determine which cards came from which choice.

A♣ A♦ A♥

A♠ K♦

A♣ A♦ A♠

A♥ K♦


A♣ A♠ A♥

A♦ K♦

A♠ A♦ A♥

A♣ K♦

Is the other
argument
correct? How do
I avoid fallacious
reasoning?



- 1) Choose 3 of 4 aces
- 2) Choose 2 non-ace cards

A♣ Q♠ A♦ A♥ K♦

Sleuth reasons:

The aces came from the first choice and the non-aces came from the second choice.

- 1) Choose 4 of 4 aces
- 2) Choose 1 non-ace

A♣ A♠ A♦ A♥ K♦

Sleuth reasons:

The aces came from the first choice and the non-ace came from the second choice.



Study Bee

- Correspondence Principle
If two finite sets can be placed into 1-1 onto correspondence, then they have the same size
- Choice Tree
- Product Rule
two conditions
- Counting by complementing
it's sometimes easier to count the "opposite" of something
- Binomial coefficient
Number of r sets of an n set