

Conjugate Gradient Method Steepest Descent

Spectral
10/27/09
v2

Goal: Solve $Ax=b$ A spd $A\bar{u}=b$

Start with steepest descent

Consider "Elliptical Bowl"

$$G(u) = (\bar{u} - u)^T A (\bar{u} - u)$$

$$\min_u G(u) = \bar{u}$$

Major axes are eigenvector of A .

note $G(u) = \bar{u}^T A \bar{u} - 2\bar{u}^T A u + u^T A u$
 $= \underbrace{\bar{u}^T b}_{\text{constant}} - 2\bar{u}^T b + u^T A u.$

Suffice min $F(u) = \frac{1}{2} u^T A u - u^T b$

Recall $\nabla F = \begin{pmatrix} \frac{dF}{du_1} \\ \vdots \\ \frac{dF}{du_n} \end{pmatrix}$

$$\nabla F = \frac{1}{2} \nabla u^T A u - \nabla u^T b$$

Claim $\frac{d u^T b}{d u_i} = b_i$ pf $\lim_{h \rightarrow 0} \frac{(u + h e_i)^T b - u^T b}{h}$

$$= \lim_{h \rightarrow 0} \frac{h e_i^T b}{h} = b_i$$

Check $\frac{d u^T A u}{d u_i} = 2(Au)_i$

$\nabla F(u) = Au - b$ residual $r = b - Au$

Note Richardson Alg

$$u^{(n+1)} = u^{(n)} + \lambda (b - A x^{(n)}) \quad \lambda = 1$$

• RA is steepest descent, step size $\lambda = 1$

Goal: Pick λ minimize $F(u^{(n)} + \lambda r)$

Set $u = u^{(n)}$

$$F(u + \lambda r) = \frac{1}{2} (u + \lambda r)^T A (u + \lambda r) - b^T (u + \lambda r)$$

$$\frac{dF(u + \lambda r)}{d\lambda} = r^T A u + \lambda r^T A r - b^T r$$

$$r^T (A u - b) + \lambda r^T A r$$

$$-r^T r + \lambda r^T A r$$

$$\frac{dF}{d\lambda} = 0$$

$$-r^T r + \lambda r^T A r = 0$$

$$\lambda = \frac{r^T r}{r^T A r}$$

Steepest Descent

Initial guess $u^{(0)}$

$$u^{(n+1)} = u^{(n)} + \lambda r$$

$$r = b - Au^{(n)}$$

$$\lambda = r^T r / r^T A r$$

Convergence Rate

(Kantorovich Lemma) B spd (real) λ_M, λ_m for B

$$\frac{(x^T B x)(x^T B^{-1} x)}{(x^T x)^2} \leq \frac{(\lambda_M + \lambda_m)^2}{4\lambda_M \lambda_m}$$

pt See SAAD Chap 5 page 132

Def $\|x\|_A = (x^T A x)^{1/2}$

$x \perp_A y$ if $x^T A y = 0$

note $x^T A y = (A^{1/2} x)^T (A^{1/2} y)$

$x \perp_A y$ iff $(A^{1/2} x) \perp A^{1/2} y$

Thm A spd, error in steepest descent

$\varepsilon^{(k)} = \bar{u} - u^{(k)}$ then

$$\|\varepsilon^{(k+1)}\|_A \leq \left(\frac{\lambda_M - \lambda_m}{\lambda_M + \lambda_m} \right) \|\varepsilon^{(k)}\|_A$$

$$= \left(\frac{\kappa - 1}{\kappa + 1} \right) \|\varepsilon^{(k)}\|_A \quad \kappa = \frac{\lambda_M}{\lambda_m}$$

$$\approx \left(1 - \frac{2}{\kappa} \right) \|\varepsilon^{(k)}\|_A$$


pf SAAD Chap 5 133 p.

Steepest Descent an example

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 10 \end{pmatrix} \quad b = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \bar{u} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad u^{(0)} = \begin{pmatrix} 9 \\ 1 \end{pmatrix}$$

$$r = b - Au^{(0)} = - \begin{pmatrix} 1 & 0 \\ 0 & 10 \end{pmatrix} \begin{pmatrix} 9 \\ 1 \end{pmatrix} = - \begin{pmatrix} 9 \\ 10 \end{pmatrix}$$

"Elliptical Bowl"



$$\lambda = \frac{r^T r}{r^T A r} = \frac{181}{81 + 1000} = \frac{181}{1081}$$

$$u^{(1)} = \begin{pmatrix} 9 \\ 1 \end{pmatrix} - \left(\frac{181}{1081} \right) \begin{pmatrix} 9 \\ 10 \end{pmatrix}$$

$$u_2^{(1)} = 1 - \frac{1810}{1081} = \frac{-729}{1081} \approx -3/4$$

We over shot!

Krylov Subspaces

$$Ax=b \quad \& \quad A\bar{x}=b \quad \text{initial guess } x^{(0)}$$

Def n th Krylov subspace

$$\mathcal{K}_n = \langle b, Ab, A^2b, \dots, A^{n-1}b \rangle$$

If $x^{(0)} \neq 0$ affine space $x^{(0)} + \mathcal{K}_n$ Krylov space

Why is \mathcal{K}_n of interest?

Assume $x^{(0)} = 0$

Claim Given an iterative method

$$x^{(n+1)} = \alpha_n x^{(n)} + \beta_n (b - Ax^{(n)}) \quad \text{then}$$

$$x^{(n)} \in \mathcal{K}_n$$

pf Induction on n

8

$n=1$ then $x^{(1)} = \beta b \in \mathcal{K}_1$

Claim If $x^{(n)} \in \mathcal{K}_n$ then $Ax^{(n)} \in \mathcal{K}_{n+1}$

Assume $x^{(n)} = a_1 b + \dots + a_{n-1} A^{n-1} b$

$$Ax^{(n)} = a_1 Ab + \dots + a_{n-1} A^n b \in \mathcal{K}_{n+1}$$

$\therefore x^{(n+1)} \in \mathcal{K}_{n+1}$

All our methods do is find $x^{(n)} \in \mathcal{K}_n$!

Best would be to find $\operatorname{Argmin}_{x^{(n)} \in \mathcal{K}_n} \|\bar{x} - x\|_2$

Not clear how to do this!

Next best thing

$$\operatorname{Argmin}_{x^{(n)} \in \mathcal{K}_n} \|\bar{x} - x^{(n)}\|_A$$

This we can do! It is called Conjugate Gradient.

Conjugate Gradient Iteration

Init^o: $x_0 = 0$ $r_0 = b$ $p_0 = r_0$

Iteration i:

$$\alpha_n = r_{n-1}^T r_{n-1} / p_{n-1}^T A p_{n-1} \quad \text{step length}$$

$$x_n = x_{n-1} + \alpha_n p_{n-1} \quad \text{approx solution}$$

$$r_n = r_{n-1} - \alpha_n A p_{n-1} \quad \text{residual}$$

$$\beta_n = r_n^T r_n / r_{n-1}^T r_{n-1} \quad \text{improvement}$$

$$p_n = r_n + \beta_n p_{n-1} \quad \text{search direction}$$

Claim $r_n = b - Ax_n$

pf induct on n $n=0$

assume true for $n-1$

$$b - Ax_n = b - A(x_{n-1} + \alpha_n p_{n-1})$$

$$= b - Ax_{n-1} - \alpha_n A p_{n-1}$$

$$= r_{n-1} - \alpha_n A p_{n-1} = r_n$$

Thm A EG for spd A for prob $Ax=b$ then while $r_{r_1} \neq 0$

$$\mathcal{K}_n = \langle X_1, \dots, X_n \rangle = \langle P_0, \dots, P_{n-1} \rangle$$

$$\langle r_0, \dots, r_{n-1} \rangle = \langle b, Ab, \dots, A^{n-1}b \rangle$$

$$\& r_n^T r_j = 0 \quad (j < n)$$

$$\& P_n^T A P_j = 0 \quad (j < n)$$

pt See Trefethen & Bau Chap 38

Thm B $A^{m \times m}$ $x = b$, A spd, CG

1) While $r_{n-1} \neq 0$ then x_n unique point in \mathcal{K}_n
minimizing $\|e_n\|_A$

2) $\|e_n\|_A \leq \|e_{n-1}\|_A$

3) $\exists n \leq m$ st $e_n = 0$

pf $x_n \in \mathcal{K}_n$ by previous thm.

let $x \in \mathcal{K}_n$ be arbitrary.

$$\Delta x = x_n - x$$

$$e = \bar{x} - x = e_n + \Delta x$$

$$\|e\|_A^2 = (e_n + \Delta x)^T A (e_n + \Delta x)$$

$$e_n^T A e_n + (\Delta x)^T A (\Delta x) + 2 e_n^T A (\Delta x)$$

note $A e_n = A(\bar{x} - x_n) = b - A x_n = r_n$

$$2 e_n^T A \Delta x = 2 r_n^T A \Delta x = 0 \quad r_n \perp_A \mathcal{K}_n$$

$$\|e\|_A^2 = e_n^T A e_n + (\Delta x)^T A (\Delta x)$$

$$A \text{ spd} \Rightarrow x \neq x_n \text{ then } \|e\|_A^2 > \|e_n\|_A^2$$

$$2) \mathcal{K}_n \subseteq \mathcal{K}_{n+1} \Rightarrow \|e_n\|_A \leq \|e_{n-1}\|_A$$

3) For some $n \leq m$ $\mathcal{K}_n = \mathbb{R}^m$ and done.

CG & Polynomial Approx

12

Prob Find poly P_n , $P_n(0)=1$ $\deg(P_n) \leq n$

minimizing $\|P_n(A)e_0\|_A$

Thm If $r_{n-1} \neq 0$ for CG then $\exists!$ P_n , $P_n(0)=1$

1) x_n has error $e_n = P_n(A)e_0$ $P_n(0)=1$

$$2) \frac{\|e_n\|_A}{\|e_0\|_A} = \inf_{P_n} \frac{\|P(A)e_0\|_A}{\|e_0\|_A} \leq \inf_{P_n} \max_{\lambda \in \lambda(A)} |P(\lambda)|$$

pf 1) See page 12A.

2) ($=$) follows from Thm B.

(\leq) Suppose $e_0 = \sum_{j=1}^m a_j v_j$ (eigen expansion of A)

$$P(A)e_0 = \sum a_j P(\lambda_j) v_j$$

pf of D)

By Thm A $X_n \in \mathcal{X}_n$ where $X_0 = 0$

thus $X_n = Q_{n-1}(A)b$ some polynomial $Q_{n-1}(z)$

$$\deg(Q_{n-1}(z)) \leq n-1$$

Consider $P_n(z) = 1 - z \cdot Q(z)$ $\deg(P_n) \leq n$

Claim $\varepsilon_n = P_n(A)\varepsilon_0$ where $\varepsilon_0 = \bar{X}$

$$P_n(A)\varepsilon_0 = (1 - A Q_{n-1}(A))\bar{X}$$

$$= \bar{X} - Q_{n-1}(A)(A\bar{X})$$

$$= \bar{X} - Q_{n-1}(A)b$$

$$= \bar{X} - X_n = \varepsilon_n$$

$$\|e_0\|_A = \sum_{j=1}^m a_j^2 \lambda_j \quad \|e_n\|_A = \sum a_j^2 p_n(\lambda_j)^2 \lambda_j$$

$$\frac{\|e_n\|_A}{\|e_0\|_A} \leq \max \frac{a_j^2 p_n(\lambda_j)^2 \lambda_j}{a_j^2 \lambda_j} = \max_j p_n(\lambda_j)$$

Thm CG to solve $Ax=b$, A spd

$$\frac{\|e_n\|_A}{\|e_0\|_A} \leq 2 \left(\frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1} \right)^n$$

pf

The Chebyshev polynomial

$$\frac{\|e_n\|_A}{\|e_0\|_A} \leq \max_{\lambda \in \lambda(A)} |P_n(A)| \leq \max_{\lambda \in \lambda(A)} |\overline{T}_n(A)|$$

$$\leq 2 \left(\frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1} \right)^n$$