

# Constructive Logic (15-317), Fall 2019

## Assignment 4: Quantification and Arithmetic

Instructor: Karl Crary  
TAs: Avery Cowan, David Kahn, Siva Somayyajula

Due: Friday, September 27, 2019, 11:59 pm

### 1 Quantification

It is important to note that quantification extends as far to the right as syntactically possible. For example, the proposition  $\exists x : \tau. A(x) \supset \forall x : \tau. A(x)$  should be interpreted as  $\exists x : \tau. (A(x) \supset \forall x : \tau. A(x))$  and not as  $(\exists x : \tau. A(x)) \supset (\forall x : \tau. A(x))$ . Tutch implements the same convention.

#### 1.1 Distributivity properties

In class, we saw that universal quantification distributes over conjunction, that is,

$$(\forall x : \tau. A(x) \wedge B(x)) \equiv (\forall x : \tau. A(x)) \wedge \forall x : \tau. B(x) \text{ true.}$$

In this section, we will explore various other distributivity properties.

**Task 1** (10 points). Dually, existential quantification distributes over disjunction, that is,

$$(\exists x : \tau. A(x) \vee B(x)) \equiv (\exists x : \tau. A(x)) \vee \exists x : \tau. B(x) \text{ true.}$$

In this task, you will show this equivalence by giving a natural deduction proof of each of the following directions:

- $(\exists x : \tau. A(x) \vee B(x)) \supset (\exists x : \tau. A(x)) \vee \exists x : \tau. B(x) \text{ true}$
- $(\exists x : \tau. A(x)) \vee (\exists x : \tau. B(x)) \supset \exists x : \tau. A(x) \vee B(x) \text{ true}$

#### 1.2 Constructive and classical quantification

**Task 2** (9 points). For each of the following judgments, give a constructive natural deduction proof and the corresponding proof term if it is constructively valid. If it is not constructively valid, state this. *N.B. The following judgments are all classically valid.*

- $(\neg \forall x : \tau. \neg A(x)) \supset \exists x : \tau. A(x) \text{ true}$
- $(\exists x : \tau. A(x)) \supset \neg \forall x : \tau. \neg A(x) \text{ true}$

### 1.3 Tutch, Quantified

Tutch uses the concrete syntax  $?x:t.A(x)$  and  $!x:t.A(x)$  for  $\exists x : \tau.A(x)$  and  $\forall x : \tau.A(x)$ , respectively. We encourage you to review the scoping rules for quantifiers described at the beginning of Section 1 of this assignment before starting this portion of the assignment. Please see the Tutch manual for more information on how to use quantifiers in Tutch.

**Task 3** (6 points). Prove each of the following propositions using Tutch. Place the proof for part a (and only the proof for part a) in `hw4_8a.tut`, ..., and the proof for part c (and only the proof for part c) in `hw4_8c.tut`.

- a. proof apply :  $(!x:t.A(x) \Rightarrow B(x)) \Rightarrow (!x:t.A(x)) \Rightarrow (!x:t.B(x));$
- b. proof instance :  $(!x:t.A(x)) \ \& \ (?y:t.B(y)) \Rightarrow ?z:t.A(z);$
- c. proof frobenius :  $(R \ \& \ ?x:t.Q(x)) \Leftrightarrow ?x:t.(R \ \& \ Q(x));$

## 2 Heyting Arithmetic

Recall that our discussion of quantified logic glossed over the judgment  $t : \tau$  assigning terms to types and its interaction with the rest of the logic. Here, we'll formally define the system of Heyting arithmetic that we discussed in class.

### 2.1 Natural Numbers

First, here are the rules for the judgment  $n : \text{nat}$ .

$$\frac{}{0 : \text{nat}} \text{nat}I_0 \qquad \frac{x : \text{nat}}{S(x) : \text{nat}} \text{nat}I_S$$

$$\frac{\frac{x : \text{nat} \quad C[0] \text{ true} \quad \frac{\frac{y : \text{nat} \quad C[y] \text{ true}}{\vdots} C[S(y)] \text{ true}}{C[x] \text{ true}}}{\text{nat}E^{y,u}}$$

### 2.2 Equality

Now, we'll add a new atomic proposition, equality, and define the truth judgment at equality.

$$\frac{}{0 = 0 \text{ true}} = I_{00} \qquad \frac{x = y \text{ true}}{S(x) = S(y) \text{ true}} = I_{SS}$$

$$\frac{0 = S(x) \text{ true}}{C \text{ true}} = E_{0S} \qquad \frac{S(x) = 0 \text{ true}}{C \text{ true}} = E_{S0} \qquad \frac{S(x) = S(y) \text{ true}}{x = y \text{ true}} = E_{SS}$$

### 2.3 Primitive Recursion

Lastly, we need the ability to define primitive recursive functions like addition and multiplication *within* the theory of the natural numbers so that we may reason about them in the natural deduction system with the induction principle introduced two subsections ago.

$$\frac{}{R(0; t_0; x, r.t_S) \Rightarrow_R t_0} \Rightarrow_R I_0 \quad \frac{}{R(S(n); t_0; x, r.t_S) \Rightarrow_R [R(n; t_0; x, r.t_S)/r][n/x]t_S} \Rightarrow_R I_S$$

$$\frac{A[x] \text{ true} \quad x \Rightarrow_R y}{A[y] \text{ true}} \Rightarrow_R E_1 \quad \frac{A[y] \text{ true} \quad x \Rightarrow_R y}{A[x] \text{ true}} \Rightarrow_R E_2$$

**Task 4** (10 points). Let  $m + n$  be defined by  $R(m; n; x, r.S(r))$ . Show that 0 is the right additive identity i.e. give a derivation of  $\forall m : \text{nat}. R(m; 0; x, r.S(r)) = m \text{ true}$ .