

Algorithms (COT 6405): Solutions 4

Problem 1

Give an example of functions $f(n)$ and $g(n)$ that satisfy all of the following conditions:

$$\begin{aligned}f(n) &= O(g(n)) \\f(n) &\neq \Theta(g(n)) \\f(n) &\neq o(g(n))\end{aligned}$$

Consider the following two functions:

$$\begin{aligned}f(n) &= 1 \\g(n) &= \begin{cases} 1 & \text{if } n \text{ is even;} \\ n & \text{if } n \text{ is odd.} \end{cases}\end{aligned}$$

Since $f(n) \leq g(n)$, we immediately conclude that $f(n) = O(g(n))$. For even n , the function $f(n)$ is of the same order as $g(n)$, which means that $f(n) \neq o(g(n))$. On the other hand, for odd n , $f(n)$ grows asymptotically slower than $g(n)$, which implies that $f(n) \neq \Theta(g(n))$.

Problem 2

Prove the following transitivity property of asymptotic bounds:

$$\text{if } f(n) = \Theta(g(n)) \text{ and } g(n) = \Theta(h(n)), \text{ then } f(n) = \Theta(h(n)).$$

Since $f(n) = \Theta(g(n))$, we conclude that there are some positive constants c_1 , c_2 , and n_1 such that, for all $n \geq n_1$, we have:

$$c_1 g(n) \leq f(n) \leq c_2 g(n).$$

Similarly, since $g(n) = \Theta(h(n))$, there are some positive constants c_3 , c_4 , and n_2 such that, for all $n \geq n_2$, we have:

$$c_3 h(n) \leq g(n) \leq c_4 h(n).$$

We combine these two inequalities as follows:

$$c_1 c_3 h(n) \leq c_1 g(n) \leq f(n) \leq c_2 g(n) \leq c_2 c_4 h(n).$$

We now define three new constants, c_5 , c_6 , and n_3 :

$$\begin{aligned}c_5 &= c_1 c_3, \\c_6 &= c_2 c_4, \\n_3 &= \max(n_1, n_2).\end{aligned}$$

Then, the last inequality implies that, for every $n \geq n_3$, we have:

$$c_5 h(n) \leq f(n) \leq c_6 h(n).$$

This inequality means that, by definition, $f(n) = \Theta(h(n))$.