

Partial Differential Equations

Paul Heckbert

Computer Science Department
Carnegie Mellon University

Differential Equation Classes 1


- dimension of unknown:
 - *ordinary differential equation* (ODE) – unknown is a function of one variable, e.g. $y(t)$
 - *partial differential equation* (PDE) – unknown is a function of multiple variables, e.g. $u(t,x,y)$
- number of equations:
 - *single* differential equation, e.g. $y'=y$
 - *system* of differential equations (*coupled*), e.g. $y_1'=y_2, y_2'=-g$
- *order*
 - n th order DE has n th derivative, and no higher, e.g. $y''=-g$

Differential Equation Classes 2

- linear & nonlinear:
 - *linear* differential equation: all terms linear in unknown and its derivatives
 - e.g.
 - $x'' + ax' + bx + c = 0$ – linear
 - $x' = t^2 x$ – linear
 - $x'' = 1/x$ – nonlinear


PDE's in Science & Engineering 1

- Laplace's Equation: $\nabla^2 u = u_{xx} + u_{yy} + u_{zz} = 0$
 - unknown: $u(x,y,z)$
 - gravitational / electrostatic potential
- Heat Equation: $u_t = a^2 \nabla^2 u$
 - unknown: $u(t,x,y,z)$
 - heat conduction
- Wave Equation: $u_{tt} = a^2 \nabla^2 u$
 - unknown: $u(t,x,y,z)$
 - wave propagation



PDE's in Science & Engineering 2

- Schrödinger Wave Equation
 - quantum mechanics
 - (electron probability densities)



- Navier-Stokes Equation
 - fluid flow (fluid velocity & pressure)



2nd Order PDE Classification

- We classify conic curve $ax^2 + bxy + cy^2 + dx + ey + f = 0$ as ellipse/parabola/hyperbola according to sign of discriminant $b^2 - 4ac$.
- Similarly we classify 2nd order PDE $au_{xx} + bu_{xy} + cu_{yy} + du_x + eu_y + fu + g = 0$:

$b^2 - 4ac < 0$ – elliptic (e.g. equilibrium)

$b^2 - 4ac = 0$ – parabolic (e.g. diffusion)

$b^2 - 4ac > 0$ – hyperbolic (e.g. wave motion)

For general PDE's, class can change from point to point

Example: Wave Equation

- $u_{tt} = c u_{xx}$ for $0 \leq x \leq 1, t \geq 0$
- initial cond.: $u(0,x) = \sin(\pi x) + x + 2, u_t(0,x) = 4\sin(2\pi x)$
- boundary cond.: $u(t,0) = 2, u(t,1) = 3$
- $c = 1$
- unknown: $u(t,x)$

- simulated using Euler's method in t

Wave Equation: Numerical Solution

$$u_j^{k+1} = 2u_j^k - u_j^{k-1} + c \frac{\Delta t^2}{\Delta x^2} (u_{j+1}^k - 2u_j^k + u_{j-1}^k)$$

`u0 = ...`

`u1 = ...`

`for t = 2*dt:dt:endt`


`u2(2:n) = 2*u1(2:n)-u0(2:n)`

`+c*(dt/dx)^2*(u1(3:n+1)-2*u1(2:n)+u1(1:n-1));`


`u0 = u1;`

`u1 = u2;`


`end`




Wave Equation Results




Wave Equation Results



Wave Equation Results



Poor results when dt too big



PDE Solution Methods

- Discretize in space, transform into system of IVP's
- Discretize in space and time, finite difference method.
- Discretize in space and time, finite element method.
 - Latter methods yield sparse systems.
- Sometimes the geometry and boundary conditions are simple (e.g. rectangular grid);
- Sometimes they're not (need mesh of triangles).