Vertex Cover on k-Uniform Hypergraphs is Hard to Approximate within Factor $(k-3-\varepsilon)$

Irit Dinur* Venkatesan Guruswami† Subhash Khot‡

April, 2002

Abstract

Given a k-uniform hypergraph, the Ek-Vertex-Cover problem is to find a minimum subset of vertices that "hits" every edge. We show that for every integer $k \geq 5$, Ek-Vertex-Cover is NP-hard to approximate within a factor of $(k-3-\varepsilon)$, for an arbitrarily small constant $\varepsilon > 0$.

This almost matches the upper bound of k for this problem, which is attained by the straightforward greedy approximation algorithm. The best previously known hardness result was due to Holmerin [Hol02a], who showed the NP-hardness of approximating Ek-Vertex-Cover within a factor of $k^{1-\varepsilon}$.

We present two constructions: one with a simple purely combinatorial analysis, showing Ek-Vertex-Cover to be NP-hard to approximate to within a factor $\Omega(k)$, followed by a stronger construction that obtains the $(k-3-\varepsilon)$ inapproximability bound. The latter construction introduces a novel way of combining ideas from Dinur and Safra's paper [DS02] and the notion of covering complexity introduced by Guruswami, Håstad and Sudan [GHS00]. This also allows us to prove a hardness factor of $(k-1-\varepsilon)$ assuming the hardness of $O(\log n)$-coloring a c-colorable graph for some fixed $c \geq 3$.

1 Introduction

Given a k-uniform hypergraph, $G = (V, E)$ with vertices V and hyperedges $E \subseteq \binom{V}{k} \stackrel{def}{=} \{e \subseteq V \mid |e| = k\}$, a vertex-cover in G is a subset $S \subseteq V$ that intersects each edge. An *independent set* in G is a subset whose complement is a vertex cover, or in other words is a subset of vertices that contains no edge. The Ek-Vertex-Cover problem consists of finding a minimum size vertex cover in a k-uniform hypergraph. This problem is alternatively called the minimum hitting set problem with sets of size k (and is equivalent to the set cover problem where each element of the universe occurs in exactly k sets).

We show that this problem is NP-hard to approximate within $(k-3-\varepsilon)$ for an arbitrarily small constant $\varepsilon > 0$. The result is almost tight as this problem is approximable to within factor k by repeatedly selecting one arbitrary hyperedge, adding all its vertices into the cover and and removing all the "covered" hyperedges. The best known algorithm [Hal00] gives only a slight improvement on this greedy algorithm, achieving an approximation factor of $k - o(1)$.

*School of Mathematics, Institute for Advanced Study, Princeton, NJ 08540. iritd@ias.edu

†University of California at Berkeley, Miller Institute for Basic Research in Science, Berkeley, CA 94720. venkat@lcs.mit.edu

‡Department of Computer Science, Princeton University, Princeton, NJ 08544. khot@cs.princeton.edu

Previous Hardness Results

This problem was suggested by Trevisan [Tre01] who initiated a study of bounded degree instances of certain combinatorial problems. There it was shown that this problem is hard to approximate within a factor of $k^{1/19}$. Holmerin [Hol02b] showed that E4-Vertex-Cover is NP-hard to approximate within $(2-\varepsilon)$, and more recently [Hol02a] that Ek-Vertex-Cover is NP-hard to approximate within $k^{1-\varepsilon}$. Goldreich [Gol01] found a simple 'FGLSS'-type [FGL$^+$91] reduction (involving no use of the long-code, a crucial component in most recent PCP constructions) to obtain a hardness factor of $(2-\varepsilon)$ for Ek-Vertex-Cover for some constant k.

Our Results

We present two constructions: one that attains a hardness factor of $\Omega(k)$ (already improving the best previously known result) with a simple, purely combinatorial analysis, and one that is stronger attaining the inapproximability factor of $(k-3-\varepsilon)$.

Our "simple" construction follows that of [Hol02b] who showed that it is NP-hard to approximate E4-Vertex-Cover to within a factor $(2-\varepsilon)$. Taking a new "set-theoretic" viewpoint on that construction, we give a purely combinatorial proof of Holmerin's theorem (in contrast to Holmerin's use of Fourier analysis), relying solely on one new Erdős-Ko-Rado (EKR) type combinatorial lemma (Lemma 2.2) that bounds the maximal size of t-intersecting families of subsets (i.e. families in which every pair of subsets intersect on at least t elements). Taking this new 'set-theoretic' viewpoint results in a direct extension of that construction to obtain an $\Omega(k)$ inapproximability factor.

Our "strong" construction, achieving a hardness factor of $k-3-\varepsilon$ involves a novel way of combining ideas from Dinur and Safra's paper [DS02] and the notion of covering complexity introduced by Guruswami, Håstad and Sudan [GHS00].

A constraint satisfaction problem is said to have covering complexity L (see [GHS00]) if it is NP-hard to distinguish between the case when the CSP has a satisfying assignment and the case when, given any L assignments, there exists a constraint that is *not* satisfied by any of these L assignments. Guruswami et al [GHS00] show hardness of coloring 2-colorable 4-uniform hypergraphs with constantly many colors. This result is equivalent to saying that the Not-All-Equal predicate on 4 binary variables has arbitrarily high covering complexity. In our paper, we need to use a (stronger) variant of the notion of covering complexity (see Definition 4.2).

We combine the covering complexity techniques with the methods from [DS02]. We borrow a powerful tool used in their paper, namely the Friedgut's Theorem about influence of variables on Boolean functions (Theorem 4.1). Assuming that for some fixed $c \geq 3$, it is hard to color an n-vertex c-colorable graph using at most $b \log n$ for every integer b, we get a predicate on 2 variables with the necessary covering complexity and this lets us prove a factor $(k-1-\varepsilon)$ hardness for the Ek-Vertex-Cover problem.

Location of the gap: All our hardness results have the gap between sizes of the vertex cover at the "right" location. Specifically, to prove a factor $(B-\varepsilon)$ hardness we show that it is hard to distinguish between k-uniform hypergraphs that have a vertex cover of weight $\frac{1}{B}+\varepsilon$ from those whose minimum vertex cover has weight at least $(1-\varepsilon)$. This result is stronger than a gap of about B achieved, for example, between vertex covers of weight $1/B^2$ and $1/B$. Put another way, our result shows that for k-uniform hypergraphs, for $k \geq 3$, there is a fixed α such that for arbitrarily small $\varepsilon > 0$, it is NP-hard to find an independent set consisting of a fraction ε of the vertices even if the hypergraph is promised to contain an independent set comprising a fraction α of vertices. We remark that such a result is not known for graphs and seems out of reach of current techniques. (The recent factor 4/3

hardness result for vertex cover on graphs due to Dinur and Safra [DS02], for example, shows that it is NP-hard to distinguish between cases when the graph has an independent set of size $n/3$ and when no independent set has more than $n/9$ vertices.)

Organization of the Paper

We begin with some preliminaries, including the starting-point PCP theorem, and some combinatorial lemmas that are used later in the analysis of the constructions. In Section 3 we present our first construction and prove a hardness of approximation factor of $\Omega(k)$. In Section 4 we present our stronger construction proving a $(k-3-\varepsilon)$ hardness factor. In Section 5 we explain how a hardness assumption for graph coloring implies a factor $(k-1-\varepsilon)$ hardness.

Subsequent Work

Following this work, Dinur, Guruswami, Khot and Regev [DGKR02] were able to *unconditionally* show a hardness of approximation factor of $(k-1-\varepsilon)$. The techniques in this paper are quite different from the ones in [DGKR02] and could be of independent interest. Moreover, the $\Omega(k)$ hardness factor is probably still the best in terms of holding for super-constant values of k.

2 Preliminaries

For a universe R, let $P(R)$ denote its power set, i.e. the family of all subsets of R. A family $\mathcal{F} \subseteq P(R)$ is called *monotone* if $F \in \mathcal{F}, F \subseteq F'$ implies $F' \in \mathcal{F}$.

For a "bias parameter" $0 < p < 1$, the weight $\mu_p(F)$ of a set F is defined as

$$\mu_p(F) \stackrel{def}{=} p^{|F|}(1-p)^{|R \setminus F|}.$$

The weight of a family $\mathcal{F} \subseteq P(R)$ is defined as

$$\mu_p(\mathcal{F}) \stackrel{def}{=} \sum_{F \in \mathcal{F}} \mu_p(F).$$

Note that the bias parameter defines a product distribution on $P(R)$, where the probability of one subset $F \in P(R)$ is determined by independently flipping a p-biased coin to determine the membership of each element of R in the subset. We denote this distribution by μ_p.

2.1 Intersecting Families of Subsets

In this section we present some combinatorial lemmas regarding intersecting families of subsets, that will be useful later.

Definition 2.1 *For a family of subsets $\mathcal{F} \subset P(R)$, let*

$$\mathcal{F} \cap \mathcal{F} \stackrel{def}{=} \{ F_1 \cap F_2 \mid F_1, F_2 \in \mathcal{F} \}.$$

Lemma 2.2 (EKR-Core) *For every $\varepsilon, \delta > 0$, there exists some $t = t(\varepsilon, \delta) > 0$ such that for every finite R and set family $\mathcal{F} \subset P(R)$, if $\mu_{\frac{1}{2}-\delta}(\mathcal{F}) > \varepsilon$, then there exists some 'core' subset $C \in \mathcal{F} \cap \mathcal{F}$ with $|C| \leq t$.*

Proof Sketch: A family of subsets is *t-intersecting* if for every $F_1, F_2 \in \mathcal{F}$, $|F_1 \cap F_2| \geq t$. The idea is that a family cannot be t-intersecting, for large (but constant) t and still retain a non-negligible size. Thus, if $\mu_{\frac{1}{2}-\delta}(\mathcal{F}) > \varepsilon$, there exists some t for which $\mathcal{F}$ is not $t+1$ intersecting, hence $\mathcal{F}$ has a 'core' subset of size t.

For the full proof (see Section A.1), we rely on the complete intersection theorem for finite sets of [AK97] that fully characterizes the maximal t-intersecting set families. ■

Proposition 2.3 *Let $p > 0$ and let $\mathcal{F} \subseteq P(R)$. Then $\mu_{p^2}(\mathcal{F} \cap \mathcal{F}) \geq (\mu_p(\mathcal{F}))^2$.*

Proof:

$$\Pr_{F \in \mu_{p^2}}[F \in \mathcal{F} \cap \mathcal{F}] = \Pr_{F_1, F_2 \in \mu_p}[F_1 \cap F_2 \in \mathcal{F} \cap \mathcal{F}] \geq \Pr_{F_1 \in \mu_p}[F_1 \in \mathcal{F}] \cdot \Pr_{F_2 \in \mu_p}[F_2 \in \mathcal{F}] = (\mu_p(\mathcal{F}))^2$$

■

Note that when $\mathcal{F}$ is monotone and defined by exactly one 'minterm', equality holds.

Proposition 2.4 *Let $p > 0$, $\mathcal{F} \subseteq P(R)$. Let $\mathcal{F}^k \stackrel{def}{=} \{F_1 \cap \cdots \cap F_k \mid F_i \in \mathcal{F}\}$. Then,*

$$\mu_{p^k}(\mathcal{F}^k) \geq (\mu_p(\mathcal{F}))^k.$$

Proof: By induction on k. ■

2.2 Starting Point - PCP

The Parallel Repetition Theorem

As is the case for many inapproximability results, we begin our reduction from Raz's parallel repetition theorem [Raz98] which is a version of the PCP theorem that is very powerful and convenient to work with. Let $\Phi = \{\varphi_1, ..., \varphi_n\}$ be a system of local-constraints over two sets of variables, denoted X and Y. Let R_X denote the range of the X-variables and R_Y the range of the Y-variables [1]. Assume each constraint $\varphi \in \Phi$ depends on exactly one $x \in X$ and one $y \in Y$, furthermore, for every value $a_x \in R_X$ assigned to x there is exactly one value $a_y \in R_Y$ to y such that the constraint φ is satisfied. Therefore, we can write each local constraint $\varphi \in \Phi$ as a function from R_X to R_Y, and use notation $\varphi_{x \to y} : R_X \to R_Y$ (this notation is borrowed from [DS02]). Furthermore, we assume that every X-variable appears in the same number of local-constraints in Φ.

Theorem 2.5 (PCP Theorem [AS98, ALM+98, Raz98]) *Let $\Phi = \{\varphi_1, ..., \varphi_n\}$ be as above. There exists a universal constant $\gamma > 0$ such that for every constant $|R_X|$, it is NP-hard to distinguish between the following two cases:*

- **YES :** *There is an assignment $A : X \cup Y \to R_X \cup R_Y$ such that all $\varphi_1, ..., \varphi_n$ are satisfied by A, i.e. $\forall \varphi_{x \to y} \in \Phi$, $\varphi_{x \to y}(A(x)) = A(y)$.*
- **NO :** *No assignment can satisfy more than a fraction $\frac{1}{|R_X|^\gamma}$ of the constraints in Φ.* ■

[1] Readers familiar with the Raz-verifier may prefer to think concretely of $R_X = [7^u]$ and $R_Y = [2^u]$ for some number u of repetitions.

3 The "Simple" Construction

In this section, we prove the factor $\Omega(k)$ hardness result for Ek-Vertex-Cover. Our construction follows that of [Hol02b] who showed that it is NP-hard to approximate E4-Vertex-Cover within factor $(2-\varepsilon)$. Taking a new viewpoint on that construction we give a purely combinatorial proof of Holmerin's theorem (in contrast to Holmerin's use of Fourier analysis), relying solely on one new Erdős-Ko-Rado (EKR) type combinatorial lemma (Lemma 2.2) that bounds the maximal size of t-intersecting families of subsets (i.e. families in which every pair of subsets intersect on at least t elements). We then show a direct extension of that construction to obtain an $\Omega(k)$ inapproximability result for Ek-Vertex-Cover.

The use of EKR-type bounds in the context of inapproximability results was initiated in [DS02] as part of a more complicated construction and analysis for proving a hardness result for approximating vertex-cover on graphs. The structure of the problem at hand allows a very modular use of EKR-type bounds, and perhaps provides a better intuition as to why they are useful. Since EKR-type bounds are known in many cases to be tight, we believe that similar such bounds may prove fruitful for obtaining improved inapproximability results for other approximation problems.

As a warmup, let us first prove

Theorem 3.1 *For any $\delta > 0$, it is NP-hard to approximate E4-Vertex-Cover to within $2-\delta$.*

This result is already known (see [Hol02b]), albeit using more complex analysis techniques.

Proof: Start with a PCP instance, as given in theorem 2.5, namely a set of local constraints $\Phi = \{\varphi_1, .., \varphi_n\}$ over variables $X \cup Y$, whose respective ranges are R_X, R_Y. For parameters, fix $t = t(\frac{\varepsilon}{2}, \delta)$, and take $|R_X| > (\frac{2t^2}{\delta})^{1/\gamma}$ where $\gamma > 0$ is the universal constant from Theorem 2.5. From Φ, we now construct a 4-uniform hypergraph whose minimum vertex cover has weight $\approx \frac{1}{2}$ or ≈ 1 depending on whether Φ is satisfiable or not.

We present a construction of a weighted hypergraph $G = \langle V, E, \Lambda \rangle$, which can then be translated into an unweighted hypergraph via a standard duplication of vertices. The vertex set of G is

$$V \stackrel{def}{=} X \times P(R_X)$$

namely for each $x \in X$ we construct a block of vertices denoted $V[x] = \{x\} \times P(R_X)$ corresponding to all possible subsets of R_X. The weight of each vertex $\langle x, F \rangle \in V$ is

$$\Lambda(\langle x, F \rangle) \stackrel{def}{=} \frac{1}{|X|} \cdot \mu_{\frac{1}{2}-\delta}(F)$$

The hyperedges are defined as follows. For every pair of local-constraints $\varphi_{x_1 \to y}, \varphi_{x_2 \to y} \in \Phi$ sharing a mutual variable $y \in Y$, we add the hyperedge $\{\langle x_1, F_1 \rangle, \langle x_1, F_1' \rangle, \langle x_2, F_2 \rangle, \langle x_2, F_2' \rangle\}$ if and only if there is no $r_1 \in F_1 \cap F_1'$ and $r_2 \in F_2 \cap F_2'$ such that $\varphi_{x_1 \to y}(r_1) = \varphi_{x_2 \to y}(r_2)$:

$$E \stackrel{def}{=} \bigcup_{\varphi_{x_1 \to y}, \varphi_{x_2 \to y} \in \Phi} \left\{ \left\{ \langle x_1, F_1 \rangle, \langle x_1, F_1' \rangle, \langle x_2, F_2 \rangle, \langle x_2, F_2' \rangle \right\} \;\middle|\; \varphi_{x_1 \to y}(F_1 \cap F_1') \cap \varphi_{x_2 \to y}(F_2 \cap F_2') = \phi \right\}$$

where the union is taken over all pairs of local-constraints with a mutual variable y.

Lemma 3.2 (Completeness) *If Φ is satisfiable, then G has a vertex cover whose weight is $\leq \frac{1}{2} + \delta$.*

Proof: Assume a satisfying assignment $A : X \cup Y \to R_X \cup R_Y$ for Φ. The following set is a vertex cover of G:

$$\{\langle x, F\rangle \in V \mid x \in X,\ A(x) \notin F\} \tag{1}$$

For every hyperedge $e = \{\langle x_1, F_1\rangle, \langle x_1, F_1'\rangle, \langle x_2, F_2\rangle, \langle x_2, F_2'\rangle\}$ either $A(x_1) \notin F_1 \cap F_1'$ or $A(x_2) \notin F_2 \cap F_2'$, otherwise since $A(x_1), A(x_2)$ agree on every mutual Y-variable, we have $\varphi_{x_1 \to y}(F_1 \cap F_1') \cap \varphi_{x_2 \to y}(F_2 \cap F_2') \neq \phi$, and e would not have been a hyperedge.

Now note that the weight of the family $\{F \mid A(x) \notin F\}$ w.r.t. the bias parameter $(\frac{1}{2} - \delta)$ is $(\frac{1}{2} + \delta)$. Hence the weight of the vertex cover in (1) is $\frac{1}{2} + \delta$. ■

Lemma 3.3 (Soundness) *If G has a vertex cover whose weight is $\leq 1 - \delta$, then Φ is satisfiable.*

Proof: Let $S \subset V$ be such a vertex cover. By an averaging argument, there must be a set $X' \subseteq X$, $|X'| \geq \frac{\delta}{2}|X|$ such that for $x \in X'$, $\Pr_{v \in_\Lambda V}[v \in S \mid v \in V[x]] \leq (1 - \frac{\delta}{2})$. For each of these blocks, define

$$\mathcal{F}_x = \{F \in P(R_X) \mid \langle x, F\rangle \notin S\}$$

It follows immediately that $\forall x \in X',\ \mu_{\frac{1}{2}-\delta}(\mathcal{F}_x) \geq \frac{\delta}{2}$. The key observation is that due to Lemma 2.2 there exists some "core" subset $C \in \mathcal{F}_x \cap \mathcal{F}_x$ whose size is $|C| \leq t = t(\frac{\delta}{2}, \delta)$. In other words, there are two subsets which we denote $F_x^1, F_x^2 \in \mathcal{F}_x$, such that $\left|F_x^1 \cap F_x^2\right| \leq t$.

We next translate these "cores" into an assignment satisfying more than $\frac{1}{|R_X|^\gamma}$ fraction of Φ. Let $x_1, x_2 \in X'$, and denote their cores respectively by $C_{x_1} = F_{x_1}^1 \cap F_{x_1}^2, C_{x_2} = F_{x_2}^1 \cap F_{x_2}^2$. The next observation is that for every $\varphi_{x_1 \to y}, \varphi_{x_2 \to y} \in \Phi$ with $x_1, x_2 \in X'$, there always exists some $r_1 \in C_{x_1}$ and $r_2 \in C_{x_2}$ such that $\varphi_{x_1 \to y}(r_1) = \varphi_{x_2 \to y}(r_2)$, or in other words, we have

$$\varphi_{x_1 \to y}(C_{x_1}) \cap \varphi_{x_2 \to y}(C_{x_2}) \neq \phi\ . \tag{2}$$

Indeed, if not, then the set

$$\left\{\langle x_1, F_{x_1}^1\rangle, \langle x_1, F_{x_1}^2\rangle, \langle x_2, F_{x_2}^1\rangle, \langle x_2, F_{x_2}^2\rangle\right\}$$

would be a hyperedge not hit by S, contradicting the assumption that S is a vertex cover.

Let $Y' \subseteq Y$ denote the set of all Y variables that participate in some local-constraint with some $x \in X'$, $Y' \stackrel{def}{=} \{y \mid \varphi_{x \to y} \in \Phi, x \in X'\}$. Associate each such $y \in Y'$, with one arbitrary $x \in X'$ with $\varphi_{x \to y} \in \Phi$, and let $C_y \stackrel{def}{=} \varphi_{x \to y}(C_x) \subset R_Y$. Now define a random assignment A by independently selecting for each $x \in X', y \in Y'$ a random value from C_x, C_y respectively. Assign the rest of the variables $(X \setminus X') \cup (Y \setminus Y')$ with any arbitrary value. To complete the proof, we prove:

Proposition 3.4

$$E_A[\#\{\varphi_{x \to y} \textit{ is satisfied by } A\}] \geq \frac{\delta}{2t^2} \cdot |\Phi|$$

Proof: We will show that for any $x \in X'$, any $\varphi_{x \to y} \in \Phi$ is satisfied by A with probability $\geq \frac{1}{t^2}$; thus the expected number of local-constrains satisfied by A is $\frac{|X'|}{|X|} \cdot \frac{1}{t^2} \geq \frac{\delta}{2t^2} \cdot |\Phi|$ (because every $x \in X$ appears in the same number of local constraints). Assume $C_y = \varphi_{x' \to y}(C_{x'})$ for some $x' \in X'$ (note that $|C_y| \leq t$). By Equation (2), we have

$$C_y \cap \varphi_{x \to y}(C_x) = \varphi_{x \to y}(C_x) \cap \varphi_{x' \to y}(C_{x'}) \neq \phi\ .$$

Therefore, there is at least one value $a_x \in C_x$ such that $\varphi_{x \to y}(a_x) \in C_y$. Since for every $x \in X'$, $|C_x| \leq t$, there is at least a $\frac{1}{t^2}$ probability of having $\varphi_{x \to y}(A(x)) = A(y)$. ■

Thus, there exists some assignment A that meets the expectation, which means it satisfies $\geq \frac{\delta}{2t^2} > \frac{1}{|R_X|^\gamma}$ of the local constraints in Φ, hence Φ is satisfiable. ■

Thus we have proven that distinguishing between the case where the minimum vertex cover of G has weight at most $(\frac{1}{2}+\delta)$ and the case where it has weight at least $(1-\delta)$, enables deciding whether Φ is satisfiable or not, and is therefore NP-hard. ■

3.1 Ek-Vertex-Cover is NP hard to Approximate Within $k/3 = \Omega(k)$.

We extend the construction above to work for any constant value of $k \geq 4$. We assume without loss of generality that k is divisible by 4, and for other values of k we can use the construction for the nearest $k' = 4m$ and add $k - k'$ distinct vertices to each edge.

The vertex set for our hypergraph is the same as in the case for $k = 4$, but the weights are different. We set

$$p = (\frac{1}{2} - \delta)^{4/k} \ , \tag{3}$$

and

$$\forall v = \langle x, F \rangle \in X \times P(R_X), \qquad \Lambda(v) \stackrel{def}{=} \frac{1}{|X|} \cdot \mu_p(F)$$

The hyperedges are as follows. For every $\varphi_{x\to y}, \varphi_{x'\to y} \in \Phi$, and every $\langle x, F_1\rangle, .., \langle x, F_{\frac{k}{2}}\rangle \in V[x]$ and $\langle x', F'_1\rangle, .., \langle x', F'_{\frac{k}{2}}\rangle \in V[x']$, we add the hyperedge $\left\{\langle x, F_1\rangle, .., \langle x, F_{\frac{k}{2}}\rangle, \langle x', F'_1\rangle, .., \langle x', F'_{\frac{k}{2}}\rangle\right\}$ to E if there is no $r \in F_1 \cap \cdots \cap F_{\frac{k}{2}}$ and $r' \in F'_1 \cap \cdots \cap F'_{\frac{k}{2}}$ with $\varphi_{x\to y}(r) = \varphi_{x'\to y}(r')$. That is,

$$E \stackrel{def}{=} \bigcup_{\varphi_{x\to y}, \varphi_{x'\to y} \in \Phi} \left\{ \left\{\langle x, F_1\rangle, .., \langle x, F_{\frac{k}{2}}\rangle, \langle x', F'_1\rangle, .., \langle x', F'_{\frac{k}{2}}\rangle\right\} \;\middle|\; \varphi_{x\to y}(\cap F_i) \cap \varphi_{x'\to y}(\cap F'_i) = \phi \right\}$$

As in the case of $k = 4$,

Lemma 3.5 (Completeness) *If Φ is satisfiable, then G has a hitting set whose weight is at most $1 - p$.*

Proof: Again we take A to be a satisfying assignment, and set $S = \bigcup_{x\in X} \{\langle x, F\rangle \mid F \not\ni A(x)\}$. The weight of S is $1 - p$. ■

The proof of soundness is also quite similar to that of Lemma 3.3, with one minor twist.

Lemma 3.6 (Soundness) *If G has a hitting set whose weight is $\leq 1 - \delta$, then Φ is satisfiable.*

Proof: Let $S \subset V$ be such a hitting set. Again we consider the set $X' \subseteq X$ for which $\forall x \in X'$, $\Pr_{v\in_\Lambda V}[v \in S \mid v \in V[x]] \leq (1 - \frac{\delta}{2})$. Again $\frac{|X'|}{|X|} \geq \frac{\delta}{2}$. For each $x \in X'$ let

$$\mathcal{F}_x = \{F \in P(R_X) \mid \langle x, F\rangle \notin S\}$$

We cannot immediately apply Lemma 2.2 to find "weak-cores" for each $\mathcal{F}_x$ because lemma 2.2 works only when $p < \frac{1}{2} - \delta$. Thus, we must first consider the family $\mathcal{F}'_x = \left\{F_1 \cap \cdots \cap F_{k/4} \;\middle|\; F_i \in \mathcal{F}_x\right\}$ whose size is guaranteed to be, by Proposition 2.4,

$$\mu_{\frac{1}{2}-\delta}(\mathcal{F}'_x) = \mu_{p^{k/4}}(\mathcal{F}'_x) \geq (\mu_p(\mathcal{F}_x))^{k/4} \geq (\delta/2)^{k/4} \ .$$

Now we can deduce the existence of a "core" subset $C_x \subset R_X$, $|C_x| \leq t = t((\delta/2)^{k/4}, \delta)$, such that there are some $F_x^1, .., F_x^{k/2} \in \mathcal{F}_x$ so that $C_x = F_x^1 \cap \cdots \cap F_x^{k/2}$. Again, observe that for every $\varphi_{x\to y}, \varphi_{x'\to y} \in \Phi$, $\varphi_{x\to y}(C_x) \cap \varphi_{x'\to y}(C_{x'}) \neq \phi$. From here we can proceed as in the proof of Lemma 3.3 above to define a random assignment that satisfies an expected fraction $\frac{\delta}{2t^2}$ of the constraints in Φ. ■

All in all, we have proven

Theorem 3.7 *It is NP-hard to approximate Ek-Vertex-Cover to within a factor $k/3$.*

Proof: For a k-uniform hypergraph, we have proven in Lemmas 3.5,3.6 that it is NP-hard to distinguish between a vertex-cover of size $1-\delta$ and $1-p = 1-(\frac{1}{2}-\delta)^{4/k} < \frac{3}{k}$. The last inequality follows from

$$(1-\frac{3}{k})^{k/4} < \frac{1}{e^{3/4}} < \frac{1}{2} - \delta$$

which implies that there is no factor $\frac{k}{3}$ approximation algorithm for vertex-cover on k-uniform hypergraphs, unless P=NP. ■

4 The "Stronger" Construction

In this section we present our stronger construction, achieving a hardness of approximation factor of $(k-3-\varepsilon)$ for an arbitrarily small $\varepsilon > 0$. This construction is quite different from the one in Section 3. Let us first introduce a couple of tools we use. First, Friedgut's Theorem about influence of variables on Boolean functions (equivalently, influence of elements on set-families) and, second, a result that follows from Holmerin's result [Hol02b] on the hardness of vertex cover in 4-uniform hypergraphs.

4.1 Friedgut's "Core" Theorem

Recall that a family $\mathcal{F} \subseteq P(R)$ is called monotone if $F \in \mathcal{F}$ and $F \subseteq F'$ implies $F' \in \mathcal{F}$. For a family $\mathcal{F}$, an element $\sigma \in R$ and a bias parameter p we define the "influence of the element on the family" as

$$\text{Influence}_p(\mathcal{F}, \sigma) \stackrel{def}{=} \Pr_{F\in\mu_p}[\text{exactly one of } F\cup\{\sigma\},\ F\setminus\{\sigma\} \text{ is in } \mathcal{F}]$$

The average sensitivity of a family is defined as the sum of the influences of all elements.

$$\text{as}_p(\mathcal{F}) \stackrel{def}{=} \sum_{\sigma\in R} \text{Influence}_p(\mathcal{F}, \sigma)$$

We will use the following theorem that can be obtained by combining Russo's Theorem and Friedgut's Theorem. This theorem essentially says that a monotone family of subsets is, in some sense, determined by a 'core', see [DS02] for details.

Theorem 4.1 ([Fri98, Rus82]) *Let p be a bias parameter, $\varepsilon, \delta > 0$ be constants and ζ be an "accuracy parameter". Let $\mathcal{F} \subseteq P(R)$ be a monotone family such that $\mu_p(\mathcal{F}) \geq \delta$. Then there exists $p' \in (p, p+\varepsilon)$ and a set $C \subseteq R$ called the "core" with the following properties :*

- *The average sensitivity of the family $\mathcal{F}$ w.r.t the bias p' is at most $\frac{1}{\varepsilon}$, i.e. $as_{p'}(\mathcal{F}) \leq \frac{1}{\varepsilon}$.*
- *The size of C is a constant that depends only on $p, \delta, \varepsilon, \zeta$.*

- *If a family $\mathcal{H} \subseteq P(R \setminus C)$ is defined as*

$$\mathcal{H} \stackrel{def}{=} \{H \mid H \subseteq R \setminus C,\ C \cup H \ \in \mathcal{F}\}\ ,$$

i.e. $\mathcal{H}$ consists of all possible extensions of the core C into set that belongs to $\mathcal{F}$, then $\mu_{p'}(\mathcal{H}) \geq 1 - \zeta$ where $\mu_{p'}(\mathcal{H})$ is the weight of the family $\mathcal{H}$ under the $\mu_{p'}$-distribution over the universe $R \setminus C$.

4.2 Hardness of 4CSP

We define a constraint satisfaction problem on 4 variables (4CSP) which captures a notion related to the notion of covering complexity introduced by [GHS00]. Our hypergraph construction will be based on the hardness of this 4CSP.

Definition 4.2 *A 4CSP $\mathcal{L} = (X, \Phi)$ over a domain D is defined as follows : X is a set of variables which take values from domain D. Every $\phi \in \Phi$ is a constraint on 4 variables (which is satisfied provided the values of the 4 variables belong to a specific subset of D^4). We define YES and NO instances of the CSP as follows.*

- **YES :** *There exists an assignment $f : X \mapsto D$ to the variables which satisfies every constraint $\phi \in \Phi$ (more formally, the values assigned by f to the 4 variables in ϕ satisfy ϕ).*
- **NO :** *For any subset of variables $Y \subseteq X$, $|Y| \geq \gamma|X|$ and for any L assignments $f_1, f_2, \ldots, f_L : Y \mapsto D$, there exists a constraint $\phi \in \Phi$ such that all the 4 variables of the constraint ϕ are contained in Y and every assignment $f_i, 1 \leq i \leq L$ fails to satisfy the constraint ϕ.*

For convenience, we say that ϕ is inside Y if all the 4 variables of the constraint ϕ are in the set Y.

Theorem 4.3 *For every integer L and every constant $\gamma > 0$, it is NP-hard to distinguish whether an instance $\mathcal{L}$ of a 4CSP over Boolean domain is a YES instance or a NO instance.*

Proof: This follows immediately from a result of Holmerin [Hol02b]. He shows that for any constant $\gamma\ ' > 0$, it is NP-hard to distinguish whether an n-vertex 4-uniform hypergraph is 2-colorable or it contains no independent set of size $\gamma' n$. Now let the vertices of the 4-uniform hypergraph be variables of a 4CSP. For every edge in the hypergraph, add a Not-All-Equal constraint on its 4 vertices. When the hypergraph is 2-colorable, it means that the 4CSP has a satisfying assignment. On the other hand, if there are L assignments that satisfy every constraint inside a set of variables of size γn, it means that this set of γn vertices can be colored properly with 2^L colors and hence there exists an independent set of size $\gamma' n = (\gamma/2^L)n$. ■

Remark : The notion of hardness between the YES and NO instances here is closely related to the notion of covering complexity introduced by [GHS00]. The notion of covering complexity requires that in the NO case, no L assignments can together satisfy all the constraints. We require an even stronger condition that no L assignments can satisfy every constraint inside a set of variables Y whose size is $\gamma|X|$.

4.3 The Construction of the Hypergraph

Let $\mathcal{B}$ be the set of all l-tuples of variables of an instance $\mathcal{L}$ of a 4CSP given by Theorem 4.3. That is

$$\mathcal{B} \stackrel{def}{=} \{(x_1, x_2, \ldots, x_l) \mid x_i \in X\}$$

An l-tuple $B \in \mathcal{B}$ will be called a "block". Let R be the set of all possible "block assignments", i.e. $R \stackrel{def}{=} D^l$ is the set of all strings of length l over the domain D. Let $P(R)$ denote the family of all subsets of R, i.e.

$$P(R) \stackrel{def}{=} \{F \mid F \subseteq R\}$$

The vertex set V of the hypergraph is defined to be

$$V \stackrel{def}{=} \mathcal{B} \times P(R) = \{(B, F) \mid B \in \mathcal{B}, F \in P(R)\}$$

The vertices will have weights. Let $p = 1 - \frac{1}{k-3} - \varepsilon$ be the "bias parameter". The weight of a vertex (B, F) is $\mu_p(F)$ where

$$\mu_p(F) \stackrel{def}{=} p^{|F|}(1-p)^{|R \setminus F|}$$

To motivate the way we define the edges of the hypergraph, assume that $f : X \mapsto D$ is an assignment that satisfies every constraint. Let $f[B]$ denote the restriction of this assignment to block B. Thus $f[B] \in R$. The edges of the hypergraph will be defined in such a way that the set of vertices $\mathcal{I}_f$

$$\mathcal{I}_f = \{(B, F) \mid B \in \mathcal{B},\ f[B] \in F\} \tag{4}$$

is an independent set.

Definition 4.4 *We say that 4 blocks (B_1, B_2, B_3, B_4) are "overlapping" if they agree on some $l-1$ coordinates and the 4 variables on the remaining coordinate form a constraint in the 4CSP. More precisely, there exist variables $x_1, x_2, \ldots, x_{l-1}$ and y_1, y_2, y_3, y_4 and an index $t, 1 \leq t \leq l$ such that*

1. $B_i = (x_1, x_2, \ldots, x_{t-1}, y_i, x_t, x_{t+1}, \ldots, x_{l-1})$ *for* $i = 1, 2, 3, 4$
2. *There is a constraint $\phi \in \Phi$ on the variables (y_1, y_2, y_3, y_4).*

Note that the tuple $(\{x_j\}_{j=1}^{l-1}, \{y_i\}_{i=1}^{4}, t, \phi)$ completely characterizes the overlapping blocks.

For a block $B = (z_1, z_2, \ldots, z_l)$ and a block assignment $\sigma \in R$, let $\sigma(z_j)$ denote the value assigned by σ to the variable z_j, which is just the j^{th} coordinate of σ. For $1 \leq j \leq l$, let $\pi_j : D^l \mapsto D^{l-1}$ be the projection operator that maps a string of length l to its substring of length $l-1$ obtained by dropping the j^{th} coordinate.

Definition 4.5 *For any overlapping blocks $(B_1, B_2, B_3.B_4)$, characterized by $(\{x_j\}_{j=1}^{l-1}, \{y_i\}_{i=1}^{4}, t, \phi)$, and block assignments $\sigma^{(i)}$ to the blocks B_is, we say that these block assignments are consistent if*

1. $\pi_t(\sigma^{(1)}) = \pi_t(\sigma^{(2)}) = \pi_t(\sigma^{(3)}) = \pi_t(\sigma^{(4)})$
2. *The values $\sigma^{(1)}(y_1), \sigma^{(2)}(y_2), \sigma^{(3)}(y_3), \sigma^{(4)}(y_4)$ satisfy the constraint ϕ.*

In short, the first condition says that the assignments $\sigma^{(i)}$ must "project" down to a common assignment to the shared $(l-1)$ coordinates, and the second condition says that the 4 values on the remaining coordinate must satisfy the constraint ϕ.

Note that if $f : X \mapsto D$ is an assignment that satisfies every constraint, and $f[B]$ is the restriction of this assignment to a block B, then for any overlapping blocks $(B_1, B_2, B_3.B_4)$, the block assignments $f[B_1], f[B_2], f[B_3], f[B_4]$ are consistent.

Definition 4.6 *For overlapping blocks (B_1, B_2, B_3, B_4), and k sets $F_1, F_2, \ldots, F_{k-3}, F^{(2)}, F^{(3)}, F^{(4)} \subseteq R$, we say that these k sets are consistent if there exist block assignments $\sigma^{(i)}$ for block B_i, $1 \leq i \leq 4$, such that*

1. $\sigma^{(1)} \in F_1 \cap F_2 \cap \ldots \cap F_{k-3}$
2. $\sigma^{(i)} \in F^{(i)}$ *for* $i = 2, 3, 4$.
3. *The assignments $\sigma^{(i)}$ are consistent as per Definition 4.5.*

Remark : Whenever we talk about consistency between sets $F_1, F_2, \ldots, F_{k-3}, F^{(2)}, F^{(3)}, F^{(4)}$, we have in mind a specific set of overlapping blocks (B_1, B_2, B_3, B_4) which we will be clear from the context.

Now we are ready to define edges of the hypergraph. For overlapping blocks (B_1, B_2, B_3, B_4), and sets $F_1, F_2, \ldots, F_{k-3}, F^{(2)}, F^{(3)}, F^{(4)}$ which are **not consistent**, we define

$$\{(B_1, F_j) | j = 1, 2, ..., k-3\} \bigcup \{(B_i, F^{(i)}) | i = 2, 3, 4\}$$

to be an edge of the hypergraph. Thus every edge contains exactly k vertices, i.e. this is a k-uniform hypergraph.

Lets verify that this way of defining edges makes sense. Suppose $f : X \mapsto D$ is an assignment that satisfies every constraint. We will show that the set $\mathcal{I}_f$ (see Equation (4)) is an independent set. As observed before, for any overlapping blocks (B_1, B_2, B_3, B_4), the block assignments $f[B_i]$ are consistent. Let

$$\{(B_1, F_j) | j = 1, 2, ..., k-3\} \bigcup \{(B_i, F^{(i)}) | i = 2, 3, 4\}$$

be any k vertices in the set $\mathcal{I}_f$. We will show that the sets

$$F_1, F_2, \ldots, F_{k-3}, F^{(2)}, F^{(3)}, F^{(4)}$$

are consistent and hence these k vertices **cannot** form an edge, thus proving that $\mathcal{I}_f$ is indeed an independent set. By definition of the set $\mathcal{I}_f$, we have $f[B_1] \in F_1 \cap F_2 \cap \ldots \cap F_{k-3}$ and $f[B_i] \in F^{(i)}$ for $i = 2, 3, 4$. Since the assignments $f[B_i]$ are consistent, taking $\sigma^{(i)} = f[B_i]$ in Definition 4.6 proves the claim.

4.4 Completeness

Lemma 4.7 *If the instance (X, Φ) as in Definition 4.2 is an YES instance, then there exists an independent set of weight $p|\mathcal{B}|$ in the hypergraph constructed above.*

Proof: We will show that if there is a global assignment $f : X \mapsto D$ that satisfies every constraint, then the hypergraph constructed in Section 4.3 has a "large" independent set. As observed in the last section, the set

$$\mathcal{I}_f = \{(B, F) \mid B \in \mathcal{B},\ f[B] \in F\ \}$$

is an independent set. The weight of this set is

$$\sum_{B\in\mathcal{B}} \sum_{F:F\subseteq R, f[B]\in F} \mu_p(F) = \sum_{B\in\mathcal{B}} p = p|\mathcal{B}|$$

Thus in the completeness case, there exists an independent set of weight $p|\mathcal{B}|$. ■

4.5 Soundness

Lemma 4.8 *If the instance (X, Φ) as in Definition 4.2 is a NO instance with parameters $L = |D|^{l-1}$ and $\gamma = \delta/4$, then the hypergraph constructed above has no independent set of weight $\delta|\mathcal{B}|$.*

Before we prove the above lemma, let us first verify that together with Lemma 4.7 it proves our main hardness result:

Theorem 4.9 *For every integer $k \geq 5$ and every $\varepsilon > 0$, the vertex cover problem on k-uniform hypergraphs is NP-hard to approximate within a factor of $(k - 3 - \varepsilon)$.*

Proof: By Lemmas 4.7 and 4.8, together with Theorem 4.3, we have a gap of $(p|\mathcal{B}|,\ \delta|\mathcal{B}|)$ in the size of the independent set which corresponds to a gap $((1-p)|\mathcal{B}|,\ (1-\delta)|\mathcal{B}|)$ in the size of the vertex cover. This is a factor $\frac{1-\delta}{1-p} = \frac{1-\delta}{\frac{1}{k-3}+\varepsilon} = k - 3 - \varepsilon'$ gap where $\varepsilon' \to 0$ as $\varepsilon, \delta \to 0$. ■

It remains to prove the soundness lemma.

Proof of Lemma 4.8: We will show that if the 4CSP instance $\mathcal{L}$ is a NO instance, then the hypergraph we constructed has no independent set of size $\delta|\mathcal{B}|$. Assume on the contrary that the hypergraph has an independent set of size $\delta|\mathcal{B}|$. Call this independent set $\mathcal{I}$. We will construct a collection of $|D|^{l-1}$ assignments to a set of variables $Y, |Y| \geq \delta|X|/4$ in the 4CSP such that every constraint inside Y is satisfied by some assignment.

For every block B, define

$$\mathcal{F}[B] \stackrel{def}{=} \{F \mid F \subseteq R,\ (B, F) \in \mathcal{I}\ \}$$

A simple averaging argument shows that for at least $\delta/2$ fraction of the blocks B, we have $\mu_p(\mathcal{F}[B]) \geq \delta/2$. Defining

$$\mathcal{B}' \stackrel{def}{=} \{B \mid B \in \mathcal{B},\ \mu_p(\mathcal{F}[B]) \geq \delta/2\}$$

we have $|\mathcal{B}'| \geq \delta|\mathcal{B}|/2$.

Lemma 4.10 *For every $B \in \mathcal{B}$, the family $\mathcal{F}[B]$ can be assumed to be a monotone family of subsets of R.*

Proof: The way we define the edges of the hypergraph, it is easy to see that if (B, F) is a vertex of an independent set then we can also add (B, F') to the independent set provided $F \subseteq F'$. Thus when the independent set is maximal, every family $\mathcal{F}[B]$ is monotone. ■

Using this lemma, for every $B \in \mathcal{B}''$, the family $\mathcal{F}[B]$ is a monotone family with $\mu_p(\mathcal{F}[B]) \geq \delta/2$. Let $\zeta > 0$ be a sufficiently small "accuracy" parameter which will be fixed later. Applying Theorem 4.1, we get

Lemma 4.11 *For every block $B \in \mathcal{B}'$, there exists a real number $p[B] \in (p, p + \frac{\varepsilon}{2})$ and a set $C[B] \subseteq R$ called the "core" with the following properties :*

- $\text{as}_{p[B]}(\mathcal{F}[B]) \ \leq \ \frac{2}{\varepsilon}$.
- *The size of* $C[B]$ *is at most* Δ_0 *which is a constant depending only on* $k, \varepsilon, \zeta, \delta$.
- *Let* $\mathcal{H}[B] \subseteq R \setminus C[B]$ *defined as*

$$\mathcal{H}[B] \stackrel{def}{=} \{H \ | H \subseteq R \setminus C[B], \ C[B] \cup H \ \in \ \mathcal{F}[B]\}$$

Then we have $\mu_{p[B]}(\mathcal{H}[B]) \geq 1 - \zeta$, *where the weight of the family* $\mathcal{H}[B]$ *is measured w.r.t. the* $p[B]$*-distribution on the universe* $R \setminus C[B]$.

4.5.1 Incorporating All Elements With Some Influence: the Extended Core

Let $\eta > 0$ be a threshold parameter which will be chosen later. For every $B \in \mathcal{B}'$, we identify a set of elements $\text{Infl}[B] \subseteq R$ that have significant influence on the family $\mathcal{F}[B]$, i.e.

$$\text{Infl}[B] = \{\sigma \in R \mid \text{Influence}_{p[B]}(\mathcal{F}[B], \sigma) \geq \eta\}$$

Since $\mathcal{F}[B]$ has average sensitivity at most $\frac{2}{\varepsilon}$ and the average sensitivity is simply the sum of influences of all the elements, it follows that the size of $\text{Infl}[B]$ is at most $\frac{2}{\eta\varepsilon}$ which is a constant. Finally define the "extended core" $\text{Ecore}[B]$ as

$$\text{Ecore}[B] \stackrel{def}{=} C[B] \ \cup \ \text{Infl}[B]$$

Clearly, the extended core has size at most $\Delta = \Delta_0 + \frac{2}{\eta\varepsilon}$.

4.5.2 The Preservation Property

Given two block assignments σ, σ', and a projection $\pi_j : D^l \mapsto D^{l-1}, 1 \leq j \leq k$, we say that the two assignments are "preserved" if $\pi_j(\sigma) \neq \pi_j(\sigma')$. Since σ, σ' differ in at least one coordinate, they will be preserved with probability $1 - \frac{1}{l}$ when a projection $\pi_j, 1 \leq j \leq l$ is picked at random.

For a block $B \in \mathcal{B}'$, say that its extended core is preserved under projection π_j if every pair of elements in the extended core is preserved. In other words, the projection operator is one-to-one on the extended core.

The extended core has size at most Δ. Choosing $l = \Delta^2$, the probability that the extended core is preserved under a random projection π_j, is at least $1 - \frac{\binom{\Delta}{2}}{l} \geq \frac{1}{2}$. Hence there exists an index $j_0, 1 \leq j_0 \leq l$, such that for at least half of the the blocks in $\mathcal{B}'$, their extended core is preserved. Assume w.l.o.g. that $j_0 = l$ and $\pi = \pi_l$ denote the projection operator which acts simply by dropping the last coordinate.

$$\mathcal{B}'' \stackrel{def}{=} \{B \mid B \in \mathcal{B}', \ \text{Ecore}[B] \text{ is preserved by } \pi\}$$

As noted, $|\mathcal{B}''| \geq |\mathcal{B}'|/2 \geq \delta|\mathcal{B}|/4$. A simple averaging argument shows that we can fix variables $x_1, x_2, \ldots, x_{l-1} \in X$ such that for at least $\delta/4$ fraction of variables y, we have $(x_1, x_2, \ldots, x_{l-1}, y) \in \mathcal{B}''$. Define

$$Y \stackrel{def}{=} \{y \mid y \in X, \ (x_1, x_2, \ldots, x_{l-1}, y) \ \in \mathcal{B}''\}$$

Thus we have $|Y| \geq \delta|X|/4$. Denote by B_y the block $(x_1, x_2, \ldots, x_{l-1}, y)$.

4.5.3 Defining Assignments

Now we are ready to define assignments to the variables in set Y so that every constraint inside Y is satisfied by some assignment. There will be one assignment $f_\tau : Y \mapsto D$ for every $\tau \in D^{l-1}$. For $\tau \in D^{l-1}$ and $\alpha \in D$, let $\tau\alpha \in R = D^l$ be the concatenated string.

The assignment $f_\tau : Y \mapsto D$ is defined as

$$f_\tau(y) = \begin{cases} \alpha & \text{if} \quad \exists\ \alpha\ \in D \text{ s.t. } \tau\alpha \in \text{Ecore}[B_y] \\ \text{undefined} & \text{otherwise} \end{cases}$$

There are two things to note here. Firstly, since the extended core is preserved, there exists at most one $\alpha \in D$ such that $\tau\alpha \in \text{Ecore}[B_y]$. Thus the definition of f_τ is unambiguous. Secondly, though the assignment f_τ is undefined for some (or even all) of the variables in Y, we will still show that for every constraint ϕ inside Y, there exists an assignment f_τ such that it satisfies the constraint ϕ. We prove this in the next section.

4.5.4 Finishing the Proof

In this section, we will show that for every constraint ϕ inside the set of variables Y, there exists an assignment f_τ that satisfies this constraint. Let ϕ be a constraint on the variables $\{y_1, y_2, y_3, y_4\}$ and consider the blocks

$$B_i = B_{y_i} = (x_1, x_2, \ldots, x_{l-1}, y_i)$$

Clearly, the blocks (B_1, B_2, B_3, B_4) are overlapping. We prove our claim in several steps.

Lemma 4.12 *There exist sets $F'_1, F'_2, \ldots, F'_{k-3} \subseteq R \setminus C[B_1]$ such that*

- $\bigcap_{j=1}^{k-3} F'_j = \phi$
- $F_j \stackrel{def}{=} C[B_1] \cup F'_j \in \mathcal{F}[B_1] \quad \text{for} \quad 1 \le j \le k-3.$

In particular, $\bigcap_{j=1}^{k-3} F_j = C[B_1]$.

Proof: From Lemma 4.11, the weight of the family $\mathcal{H}[B_1]$ w.r.t. the bias parameter $p[B]$ is at least $1 - \zeta$. Noting that $p[B] \le 1 - \frac{1}{k-3} - \frac{\varepsilon}{2}$ and applying Lemma A.4, there exist sets $F'_j \subseteq R \setminus C[B_1], 1 \le j \le k-3$ whose intersection is empty. By definition of the family $\mathcal{H}[B_1]$, the sets $F_j \stackrel{def}{=} C[B_1] \cup F'_j \in \mathcal{F}[B_1]$. ■

Define

$$S \stackrel{def}{=} \{\sigma \in R \mid \text{there exists } \sigma' \in C[B_1] \text{ such that } \pi(\sigma) = \pi(\sigma')\}$$

That is, S is the set of all strings which share a common prefix of length $l-1$ with *some* string in $C[B_1]$. Clearly $|S| = |D| \cdot |C[B_1]| \le |D| \cdot \Delta_0$. For $i = 2, 3, 4$ define

$$T_i \stackrel{def}{=} S \setminus \text{Ecore}[B_i]$$

Thus T_i is a set of size at most $|D| \cdot \Delta_0$. By definition of the extended core (at the end of the Section 4.5), all elements of the set T_i have influence at most η on the family $\mathcal{F}[B_i]$ w.r.t. bias $p[B_i]$. Applying Lemma A.5, if η is small enough, there exists a set $F^{(i)} \in \mathcal{F}[B_i]$ such that $F^{(i)} \cap T_i = \phi$.

Now consider the following vertices of the hypergraph :

$$\{(B_1, F_j) \mid 1 \le j \le k-3\} \bigcup \{(B_i, F^{(i)}) \mid i = 2, 3, 4\}$$

There vertices are in the independent set $\mathcal{I}$. Hence there **cannot** be an edge on these vertices. Therefore the sets $F_1, \ldots, F_j, F^{(2)}, F^{(3)}, F^{(4)}$ are consistent (see Definition 4.6). This means that there exist block assignments $\sigma^{(i)} \in R$ such that

- $\sigma^{(1)} \in \bigcap_{j=1}^{k-3} F_j = C[B_1]$ (by Lemma 4.12).
- For $i = 2, 3, 4$, $\sigma^{(i)} \in F^{(i)}$.
- The block assignments $\sigma^{(i)}$ have the same prefix of length $l-1$, i.e. there is a string $\tau \in D^{l-1}$, and values $\alpha_i \in D$ such that $\sigma^{(i)} = \tau\alpha_i$.
- The values $(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$ satisfy the constraint ϕ.

Lemma 4.13 $\sigma^{(i)} \in \text{Ecore}[B_i]$ for $i = 1, 2, 3, 4$.

Proof: We have $\sigma^{(1)} \in C[B_1] \subseteq \text{Ecore}[B_1]$. Now consider $i = 2, 3, 4$. Since $\sigma^{(i)}$ has the same $(l-1)$-prefix with $\sigma^{(1)}$, by definition of the set S, $\sigma^{(i)} \in S$. Also $\sigma^{(i)} \in F^{(i)}$ and $F^{(i)} \bigcap T_i = \phi$. Therefore $\sigma^{(i)} \in \text{Ecore}[B_i]$. ■

From this lemma, and the way the assignment f_τ is defined, we have $f_\tau(y_i) = \alpha_i$ for $i = 1, 2, 3, 4$. Since the values $(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$ satisfy the constraint ϕ, it follows that the assignment f_τ satisfies the constraint ϕ. This finishes the proof of Lemma 4.8.

5 Improved Result Assuming Hardness of Graph Coloring

A closer inspection of our proof in the previous section shows that the slack of 3 in our hardness result comes from the fact that CSP we started with had constraints that depend on 4 variables. Consequently, we needed to split the k-sets (that comprise a hyperedge) amongst four blocks, and had to get a small core as the intersection of $(k-3)$ sets belonging to any set family with substantial weight under μ_p. This in turn limits the bias parameter p to be at most $(1 - \frac{1}{k-3})$, leading to a factor $(k-3-\varepsilon)$ hardness. If we had a 2CSP (where each constraint depends only on two variables) for which a hardness similar to Theorem 4.3 holds, then we will be able to get a hardness of approximation factor of $(k-1-\varepsilon)$. This is because we will only need a small core as the intersection of $(k-1)$ sets, and can therefore pick the bias parameter p to be $(1 - \frac{1}{k-1} - \varepsilon)$. The rest of the analysis remains unchanged.

Unfortunately, we still seem to be quite far from proving a result like Theorem 4.3 for 2CSPs. However, such a result follows if a strong hardness assumption on approximate graph coloring holds. The following lemma makes formal this connection. The proof is straightforward and uses ideas similar to that of Theorem 4.3.

Lemma 5.1 *Suppose that there exists a $c \geq 3$ such that for every positive integer b it is NP-hard to $(b \log n)$-color a c-colorable graph on n-vertices. Then there exists $d \geq 2$ such that for every integer L and every constant $\gamma > 0$, given a 2CSP over domain size d, no polynomial time algorithm can tell if it is an YES instance or a NO instance unless* P = NP *(where YES and NO instances are defined for a particular L, γ as in Definition 4.2).*

We note that the above hardness follows by a Turing reduction (unlike the many-one reductions presented in the rest of the paper). We therefore have a strong hardness result for Ek-Vertex-Cover based on the hardness assumption for graph coloring.

Theorem 5.2 *Suppose that there exists a $c \geq 3$ such that for every positive integer b it is NP-hard to $(b \log n)$-color a c-colorable graph on n-vertices. Then, for every integer $k \geq 3$ and $\varepsilon > 0$, there is no polynomial time factor $(k - 1 - \varepsilon)$ algorithm for the vertex cover problem on k-uniform hypergraphs unless* P = NP.

6 Future Work

The vertex cover in every k-uniform hypergraph can be approximated to within factor $k - o(1)$, [Hal00]. An obvious open problem is that of improving our $(k - 3 - \varepsilon)$ bound and obtaining an 'optimal' inapproximability factor of $(k - \varepsilon)$, i.e. proving NP-hardness of approximating Ek-Vertex-Cover within a factor of $(k - \varepsilon)$ for any constant $\varepsilon > 0$. This problem is especially interesting for small values of k, as the $k = 2$ case (i.e. Vertex-Cover on graphs) has received a good deal of attention yet leaving the factor $(2 - \varepsilon)$ hardness result still out of reach.

Following this work, Dinur, Guruswami, Khot and Regev [DGKR02] were able to improve our work and show a hardness-of-approximation factor of $(k - 1 - \varepsilon)$.

Another possible direction is to extend these results for larger values of k. The largest plausible value of k is $\ln n$ since the greedy set-cover algorithm can always be used to achieve a $(\ln n + 1)$ approximation on any hypergraph (here n is the number of edges in the hypergraph). Our hardness result from Section 3 gives an $\Omega(k)$ inapproximability factor (assuming NP $\not\subseteq$ DTIME($2^{\log^{O(1)} n}$)) for k up to $\log^{\gamma} n$ for some absolute constant $\gamma > 0$. We conclude with the following conjecture:

Conjecture 6.1 *It is NP-hard to approximate Ek-Vertex-Cover to within $k \cdot (1 - \varepsilon)$ for any $k \leq \ln n$ and any constant $\varepsilon > 0$.*

References

[AK97] Rudolf Ahlswede and Levon H. Khachatrian. The complete intersection theorem for systems of finite sets. *European J. Combin.*, 18(2):125–136, 1997.

[ALM+98] Sanjeev Arora, Carsten Lund, Rajeev Motwani, Madhu Sudan, and Mario Szegedy. Proof verification and the hardness of approximation problems. *Journal of the ACM*, 45(3):501–555, May 1998.

[AS98] Sanjeev Arora and Shmuel Safra. Probabilistic checking of proofs: a new characterization of NP. *Journal of the ACM*, 45(1):70–122, January 1998.

[DGKR02] Irit Dinur, Venkatesan Guruswami, Subhash Khot, and Oded Regev. Manuscript in preparation, 2002.

[DS02] Irit Dinur and Shmuel Safra. On the importance of being biased. In *Proc. 34th ACM Symp. on Theory of Computing*, 2002.

[FGL+91] Uriel Feige, Shafi Goldwasser, Láslzo Lovász, Shmuel Safra, and Mario Szegedy. Approximating clique is almost NP-complete. In *Proc. 32nd IEEE Symp. on Foundations of Computer Science*, pages 2–12, 1991.

[Fri98] Ehud Friedgut. Boolean functions with low average sensitivity depend on few coordinates. *Combinatorica*, 18(1):27–35, 1998.

[GHS00] Venkatesan Guruswami, Johan Håstad, and Madhu Sudan. Hardness of approximate hypergraph coloring. In *Proc. 41st IEEE Symp. on Foundations of Computer Science*, pages 149–158. IEEE Computer Society Press, 2000.

[Gol01] Oded Goldreich. Using the FGLSS-reduction to prove inapproximability results for minimum vertex cover in hypergraphs. ECCC Technical Report TR01-102, 2001.

[Hal00] Eran Halperin. Improved approximation algorithms for the vertex cover problem in graphs and hypergraphs. In *Proceedings of the Eleventh Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 329–337, N.Y., January 9–11 2000. ACM Press.

[Hol02a] Jonas Holmerin. Improved inapproximability results for vertex cover on k-regular hypergraphs. *Proceedings of the 29th International Colloquium on Automata, Languages, and Programming (ICALP)*, July 2002.

[Hol02b] Jonas Holmerin. Vertex cover on 4-regular hypergraphs is hard to approximate within 2 - ε. In *Proc. 34th ACM Symp. on Theory of Computing*, 2002.

[Raz98] Ran Raz. A parallel repetition theorem. *SIAM Journal on Computing*, 27(3):763–803, June 1998.

[Rus82] Lucio Russo. An approximate zero-one law. *Z. Wahrsch. Verw. Gebiete*, 61(1):129–139, 1982.

[Tre01] Luca Trevisan. Non-approximability results for optimization problems on bounded degree instances. In *Proc. 33rd ACM Symp. on Theory of Computing*, 2001.

A Some Useful Lemmas

A.1 Combinatorial Core

Lemma 2.2 (Combinatorial Core) *For every $\varepsilon, \delta > 0$, there exists some $t = t(\varepsilon, \delta) > 0$ such that for every finite R and $\mathcal{F} \subset P(R)$, if $\mu_{\frac{1}{2}-\delta}(\mathcal{F}) > \varepsilon$, then there exists some 'core' subset $C \in \mathcal{F} \cap \mathcal{F}$ with $|C| \leq t$.*

Proof: We begin by stating a continuous variant of the complete intersection theorem of Ahlswede and Khachatrian. This was already proven in [DS02] for $t = 2$ and the extension for larger t is straightforward.

Define for every $i \geq 0$, $t > 0$ and $n \geq 2i + t$,

$$\mathcal{A}^n_{i,t} \stackrel{def}{=} \{ F \in P([n]) \,|\, |F \cap [1, t+2i]| \geq t + i \}.$$

Clearly, for any $n' > n \geq 2i + t$, $\mu_p(\mathcal{A}^{n'}_{i,t}) = \mu_p(\mathcal{A}^n_{i,t})$. Denoting $\binom{[n]}{k} \stackrel{def}{=} \{F \subset [n] \,|\, |F| = k\}$, the complete intersection theorem of Ahlswede and Khachatrian states that

Theorem A.1 ([AK97]) *Let $\mathcal{F} \subseteq \binom{[n]}{k}$ be t-intersecting (i.e. for every $F_1, F_2 \in \mathcal{F}$, $|F_1 \cap F_2| \geq t$). Then,*

$$|\mathcal{F}| \leq \max_{0 \leq i \leq \frac{n-t}{2}} \left| \mathcal{A}^n_{i,t} \cap \binom{[n]}{k} \right|$$

■

The following lemma is a continuous variant of the above theorem,

Lemma A.2 *[DS02] Let $\mathcal{F} \subset P([n])$ be t-intersecting. For any $p < \frac{1}{2}$,*

$$\mu_p(\mathcal{F}) \leq \max_i \left\{\mu_p(\mathcal{A}^n_{i,t})\right\}$$

■

We define for every $t > 0$ and $p < \frac{1}{2}$, let $a_{p,t} \stackrel{def}{=} \max_i(\mu_p(\mathcal{A}^n_{i,t}))$. In order to prove Lemma 2.2 it suffices to prove that for a fixed $p < \frac{1}{2}$, $\limsup_{t\to\infty} a_{p,t} = 0$.

Note that $\mathcal{A}^n_{i,t} \subseteq \{F \in P([n]) \mid |F \cap [1, t+2i]| \geq (t+2i)/2\}$. Define $\mathcal{F}_{i,t}$ to be the family $\{F \in P([t+2i]) \mid |F| \geq (t+2i)/2\}$. We then have

$$\mu_p(\mathcal{A}^n_{i,t}) \leq \mu_p(\mathcal{F}_{i,t}) \ . \tag{5}$$

Now sets in $\mathcal{F}_{i,t}$ contain at least a fraction $1/2$ of the universe $[t+2i]$, while a random set drawn according to the product distribution μ_p has an expected fraction $p < 1/2$ of elements. By standard Chernoff bounds the probability of a set picked according to μ_p landing in $\mathcal{F}_{i,t}$ is exponentially small in t and thus tends to zero as $t \to \infty$. Hence $\mu_p(\mathcal{F}_{i,t})$ tends to zero as $t \to \infty$ (for every i). Together with (5) this shows that for each fixed $p < 1/2$, $\limsup_{t\to\infty} a_{p,t} = 0$.

■

A.2 k-wise Intersecting Families

We will use the following theorem of Frankl.

Theorem A.3 *Let $\mathcal{F} \subseteq P(R)$ where $|R| = n$ and every set in the family $\mathcal{F}$ has size m. Assume that every k sets in the family have nonempty intersection and $n > mk/(k-1)$. Then*

$$|\mathcal{F}| \leq \binom{n-1}{m-1}$$

Note that a family of sets of size m containing one fixed element has size $\binom{n-1}{m-1}$. We will use the above theorem to prove :

Lemma A.4 *Let $\varepsilon > 0$ be an arbitrarily small constant, $k \geq 2$ an integer and $p = 1 - \frac{1}{k} - \varepsilon$. Let $\mathcal{F} \subseteq P(R)$ be a family such that every k sets in this family have a nonempty intersection. Then*

$$\mu_p(\mathcal{F}) < p + \varepsilon$$

provided the universe R is sufficiently large.

Proof: Let $n = |R|$ be the size of the universe. Partition the family $\mathcal{F}$ according to different set-sizes.

$$\mathcal{F}_i \stackrel{def}{=} \{F \mid F \in \mathcal{F},\ |F| = i\}$$

With the bias parameter p, the total weight of all sets of size more than $(p + \frac{\varepsilon}{2})n$ is at most $\frac{\varepsilon}{2}$ when the universe is large enough. Hence

$$\mu_p(\mathcal{F}) \ \leq \ \frac{\varepsilon}{2} + \sum_{m \leq (p+\frac{\varepsilon}{2})n} \mu_p(\mathcal{F}_m)$$

For $m \leq (p + \frac{\varepsilon}{2})n$, we have $n > mk/(k-1)$. Since every k sets in the family $\mathcal{F}_m$ have a nonempty intersection, applying Frankl's Theorem, we get

$$|\mathcal{F}_m| \ \leq \ \binom{n-1}{m-1}$$

Noting that every set in $\mathcal{F}_m$ has weight $p^m(1-p)^{n-m}$ we have

$$\begin{aligned}
\mu_p(\mathcal{F}) &\leq \frac{\varepsilon}{2} + \sum_{m \leq (p+\frac{\varepsilon}{2})n} \binom{n-1}{m-1} p^m (1-p)^{n-m} \\
&\leq \frac{\varepsilon}{2} + p\left(\sum_m \binom{n-1}{m-1} p^{m-1} (1-p)^{(n-1)-(m-1)} \right) \\
&= \frac{\varepsilon}{2} + p
\end{aligned}$$

■

A.3 Very Small Influence

The following lemma can be found in [DS02].

Lemma A.5 *Let $\mathcal{F} \subseteq P(R)$ be a monotone family. Let T be a set of elements such that for every element $\sigma \in T$,* $\text{Influence}_p(\mathcal{F}, \sigma) < \eta$. *Assume η is small enough so that*

$$|T| \cdot \eta \cdot p^{-|T|} < \mu_p(\mathcal{F})$$

Then there exists a set $F \in \mathcal{F}$ such that $F \cap T = \phi$.