


15-251

Great Theoretical Ideas in Computer Science

Counting I: One-To-One Correspondence and Choice Trees

Lecture 6 (September 10, 2009)



If I have 14 teeth on the top and
12 teeth on the bottom, how many
teeth do I have in all?



Addition Rule

Let A and B be two disjoint finite sets

$$|A \cup B| = |A| + |B|$$

Addition of Multiple Disjoint Sets:

Let $A_1, A_2, A_3, \dots, A_n$ be disjoint, finite
sets:

$$\left| \bigcup_{i=1}^n A_i \right| = \sum_{i=1}^n |A_i|$$

Addition Rule (2 Possibly Overlapping Sets)

Let A and B be two finite sets:

$$|A \cup B| = |A| + |B| - |A \cap B|$$

Inclusion-Exclusion

If A, B, C are three finite sets,
what is the size of $(A \cup B \cup C)$?

$$|A| + |B| + |C| \\ - |A \cap B| - |A \cap C| - |B \cap C| \\ + |A \cap B \cap C|$$

Inclusion-Exclusion

If A_1, A_2, \dots, A_n are n finite sets,
what is the size of $(A_1 \cup A_2 \cup \dots \cup A_n)$?

$$\sum_i |A_i| \\ - \sum_{i < j} |A_i \cap A_j| \\ + \sum_{i < j < k} |A_i \cap A_j \cap A_k| \\ \dots \\ + (-1)^{n-1} |A_1 \cap A_2 \cap \dots \cap A_n|$$


Partition Method

To count the elements of a finite set S,
partition the elements into
non-overlapping subsets $A_1, A_2, A_3, \dots, A_n$.

$$\left| \bigcup_{i=1}^n A_i \right| = \sum_{i=1}^n |A_i|$$

Partition Method

S = all possible outcomes of one
white die and one black die.



Partition Method

S = all possible outcomes of one
white die and one black die.


Partition S into 6 sets:

- A_1 = the set of outcomes where the white die is 1.
- A_2 = the set of outcomes where the white die is 2.
- A_3 = the set of outcomes where the white die is 3.
- A_4 = the set of outcomes where the white die is 4.
- A_5 = the set of outcomes where the white die is 5.
- A_6 = the set of outcomes where the white die is 6.

Each of 6 disjoint set have size 6 = 36 outcomes

Partition Method

S = all possible outcomes where
the white die and the black die
have different values



S \equiv Set of all outcomes where the dice show different values. $|S| = ?$

$A_i \equiv$ set of outcomes where black die says i and the white die says something else.

$$|S| = \sum_{i=1}^6 |A_i| = \sum_{i=1}^6 5 = 30$$

S \equiv Set of all outcomes where the dice show different values. $|S| = ?$

B \equiv set of outcomes where dice agree.

$$|S \cup B| = \# \text{ of outcomes} = 36$$

$$|S| + |B| = 36$$

$$|B| = 6$$

$$|S| = 36 - 6 = 30$$

Difference Method

To count the elements of a finite set S , find two sets A and B such that

S and B are disjoint

and

$$S \cup B = A$$

$$\text{then } |S| = |A| - |B|$$

S \equiv Set of all outcomes where the black die shows a smaller number than the white die. $|S| = ?$

$A_i \equiv$ set of outcomes where the black die says i and the white die says something larger.

$$S = A_1 \cup A_2 \cup A_3 \cup A_4 \cup A_5 \cup A_6$$

$$|S| = 5 + 4 + 3 + 2 + 1 + 0 = 15$$

S \equiv Set of all outcomes where the black die shows a smaller number than the white die. $|S| = ?$


L \equiv set of all outcomes where the black die shows a larger number than the white die.

$$|S| + |L| = 30$$

It is clear by symmetry that $|S| = |L|$.


Therefore $|S| = 15$

“It is clear by symmetry that $|S| = |L|$?”



Pinning Down the Idea of Symmetry by Exhibiting a Correspondence

Put each outcome in S in correspondence with an outcome in L by swapping color of the dice.



Each outcome in S gets matched with exactly one outcome in L, with none left over.

$$\text{Thus: } |S| = |L|$$


Let $f : A \rightarrow B$ Be a Function From a Set A to a Set B

f is injective if and only if


$$\forall x, y \in A, x \neq y \Rightarrow f(x) \neq f(y)$$

f is surjective if and only if


$$\forall z \in B \exists x \in A f(x) = z$$




Let's Restrict Our Attention to Finite Sets




$$\exists \text{ injective (1-1) } f : A \rightarrow B \Rightarrow |A| \leq |B|$$



$$\exists \text{ surjective (onto) } f : A \rightarrow B \Rightarrow |A| \geq |B|$$



$$\exists \text{ bijective } f : A \rightarrow B \Rightarrow |A| = |B|$$




bijective f means the
inverse f^{-1} is well-defined

$$\exists \text{ bijective } f : A \rightarrow B \Rightarrow |A| = |B|$$

Correspondence Principle

If two finite sets can be placed into bijection, then they have the same size



It's one of the most important mathematical ideas of all time!

Question: How many n -bit sequences are there?

000000	\leftrightarrow	0
000001	\leftrightarrow	1
000010	\leftrightarrow	2
000011	\leftrightarrow	3
:	:	:
111111	\leftrightarrow	$2^n - 1$

Each sequence corresponds to a unique number from 0 to $2^n - 1$. Hence 2^n sequences.

$S = \{a, b, c, d, e\}$ has Many Subsets

$\{a\}, \{a, b\}, \{a, d, e\}, \{a, b, c, d, e\}, \{e\}, \emptyset, \dots$

The entire set and the empty set are subsets with all the rights and privileges pertaining thereto

Question: How Many Subsets Can Be Made From The Elements of a 5-Element Set?

a	b	c	d	e
0	1	1	0	1

$\{ \quad b \quad c \quad \quad e \}$ 1 means "TAKE IT"
0 means "LEAVE IT"

Each subset corresponds to a 5-bit sequence (using the "take it or leave it" code)

$S = \{a_1, a_2, a_3, \dots, a_n\}$, $T =$ all subsets of S
 $B =$ set of all n -bit strings

a_1	a_2	a_3	a_4	a_5
b_1	b_2	b_3	b_4	b_5

For bit string $b = b_1 b_2 b_3 \dots b_n$, let $f(b) = \{a_i \mid b_i = 1\}$

Claim: f is injective

Any two distinct binary sequences b and b' have a position i at which they differ

Hence, $f(b)$ is not equal to $f(b')$ because they disagree on element a_i

$S = \{a_1, a_2, a_3, \dots, a_n\}$, $T =$ all subsets of S
 $B =$ set of all n -bit strings

a_1	a_2	a_3	a_4	a_5
b_1	b_2	b_3	b_4	b_5

For bit string $b = b_1 b_2 b_3 \dots b_n$, let $f(b) = \{a_i \mid b_i = 1\}$


Claim: f is surjective

Let X be a subset of $\{a_1, \dots, a_n\}$.

Define $b_k = 1$ if a_k in X and $b_k = 0$ otherwise.

Note that $f(b_1 b_2 \dots b_n) = X$.


The number
of subsets of
an n-element
set is 2^n



Let $f : A \rightarrow B$ Be a Function From
Set A to Set B


f is a 1 to 1 correspondence (bijection) iff
 $\forall z \in B \exists$ exactly one $x \in A$ such that $f(x) = z$

f is a k to 1 correspondence iff
 $\forall z \in B \exists$ exactly k $x \in A$ such that $f(x) = z$




3 to 1 function



To count the number of horses in
a barn, we can count the number
of hoofs and then divide by 4



If a finite set A
has a k-to-1
correspondence
to finite set B,
then $|B| = |A|/k$



I own 3 beanies and 2
ties. How many different
ways can I dress up in a
beanie and a tie?



A Restaurant Has a Menu With 5 Appetizers, 6 Entrees, 3 Salads, and 7 Desserts

How many items on the menu?

$$5 + 6 + 3 + 7 = 21$$

How many ways to choose a complete meal?

$$5 \times 6 \times 3 \times 7 = 630$$

How many ways to order a meal if I am
allowed to skip some (or all) of the courses?

$$6 \times 7 \times 4 \times 8 = 1344$$


Leaf Counting Lemma

Let T be a depth- n tree when each node at
depth $0 \leq i \leq n-1$ has P_{i+1} children


The number of leaves of T is given by:

$$P_1 P_2 \dots P_n$$

Choice Tree



A choice tree is a rooted, directed tree with
an object called a “choice” associated with
each edge and a label on each leaf



A choice tree provides a “choice tree
representation” of a set S , if

1. Each leaf label is in S , and each
element of S is some leaf label
2. No two leaf labels are the same

We will now
combine the
correspondence
principle with the
leaf counting
lemma to make a
powerful counting
rule for choice tree
representation.

Product Rule

Suppose every object of a set S can be
constructed by a sequence of choices with P_1
possibilities for the first choice, P_2 for the
second, and so on.

IF 1. Each sequence of choices
constructs an object of type S

AND

2. No two different sequences create the
same object

THEN

There are $P_1 P_2 P_3 \dots P_n$ objects of type S

How Many Different Orderings of Deck With 52 Cards?

What object are we making? Ordering of a deck

Construct an ordering of a deck by a sequence of 52 choices:

52 possible choices for the first card;

51 possible choices for the second card;

⋮

1 possible choice for the 52nd card.

By product rule: $52 \times 51 \times 50 \times \dots \times 2 \times 1 = 52!$

A permutation or arrangement of n objects is an ordering of the objects


The number of permutations of n distinct objects is $n!$



How many sequences of 7 letters are there?

$$26^7$$

(26 choices for each of the 7 positions)




How many sequences of 7 letters contain at least two of the same letter?

$$26^7 - 26 \times 25 \times 24 \times 23 \times 22 \times 21 \times 20$$

number of sequences containing all different letters

The "Difference Principle"

Sometimes it is easiest to count the number of objects with property Q, by counting the number of objects that do not have property Q.



If 10 horses race, how many orderings of the top three finishers are there?

$$10 \times 9 \times 8 = 720$$

Number of ways of ordering, permuting, or arranging r out of n objects

n choices for first place, $n-1$ choices for second place, . . .

$$n \times (n-1) \times (n-2) \times \dots \times (n-(r-1))$$

$$= \frac{n!}{(n-r)!}$$

Ordered Versus Unordered

From a deck of 52 cards how many ordered pairs can be formed?

$$52 \times 51$$

How many unordered pairs?

$$(52 \times 51) / 2 \leftarrow \text{divide by overcount}$$

Each unordered pair is listed twice on a list of the ordered pairs

Ordered Versus Unordered

From a deck of 52 cards how many ordered pairs can be formed?

$$52 \times 51$$

How many unordered pairs?

$$(52 \times 51) / 2 \leftarrow \text{divide by overcount}$$

We have a 2-1 map from ordered pairs to unordered pairs.

Hence #unordered pairs = (#ordered pairs)/2

Ordered Versus Unordered

How many ordered 5 card sequences can be formed from a 52-card deck?

$$52 \times 51 \times 50 \times 49 \times 48$$

How many orderings of 5 cards?

$$5!$$

How many unordered 5 card hands?

$$(52 \times 51 \times 50 \times 49 \times 48) / 5! = 2,598,960$$

A combination or choice of r out of n objects is an (unordered) set of r of the n objects


The number of r combinations of n objects:

$$\frac{n!}{r!(n-r)!} = \binom{n}{r}$$

n "choose" r

The number of subsets of size r that can be formed from an n -element set is:

$$\frac{n!}{r!(n-r)!} = \binom{n}{r}$$



How Many 8-Bit Sequences Have 2 0's and 6 1's?

Tempting, but incorrect:
8 ways to place first 0, times
7 ways to place second 0

Violates condition 2 of product rule!

Choosing position i for the first 0 and then position j for the second 0 gives same sequence as choosing position j for the first 0 and then position i for the second 0

2 ways of generating the same object!

How Many 8-Bit Sequences Have 2 0's and 6 1's?

1. Choose the set of 2 positions to put the 0's. The 1's are forced.

$$\binom{8}{2}$$

2. Choose the set of 6 positions to put the 1's. The 0's are forced.

$$\binom{8}{6}$$

Symmetry In The Formula


$$\binom{n}{r} = \frac{n!}{r!(n-r)!} = \binom{n}{n-r}$$

"# of ways to pick r out of n elements"

=

"# of ways to choose the $(n-r)$ elements to omit"

Counting Cards



How Many 5-card hands Have at Least 3 As?

How Many Hands Have at Least 3 As?

$$\binom{4}{3}$$

= ways of picking 3 out of 4 aces

$$\binom{49}{2}$$

= ways of picking 2 cards out of the remaining 49 cards

$$4 \times 1176 = 4704$$

How Many Hands Have at Least 3 As?

How many hands have exactly 3 aces?


$$\begin{array}{r} \left(\begin{array}{c} 4 \\ 3 \end{array} \right) = \text{ways of picking 3 out of 4 aces} \\ \left(\begin{array}{c} 48 \\ 2 \end{array} \right) = \text{ways of picking 2 cards out of the 48 non-ace cards} \end{array} \quad \begin{array}{r} 4 \\ \times 1128 \\ \hline 4512 \end{array}$$

How many hands have exactly 4 aces?

$$\begin{array}{r} \left(\begin{array}{c} 4 \\ 4 \end{array} \right) = \text{ways of picking 4 out of 4 aces} \\ \left(\begin{array}{c} 48 \\ 1 \end{array} \right) = \text{ways of picking 1 cards out of the 48 non-ace cards} \end{array} \quad \begin{array}{r} \\ + 48 \\ \hline 4560 \end{array}$$

4704 \neq 4560

At least one of the two counting arguments is not correct!




Four Different Sequences of Choices Produce the Same Hand

$$\begin{array}{l} \left(\begin{array}{c} 4 \\ 3 \end{array} \right) = 4 \text{ ways of picking 3 out of 4 aces} \\ \left(\begin{array}{c} 49 \\ 2 \end{array} \right) = 1176 \text{ ways of picking 2 cards out of the remaining 49 cards} \end{array}$$


A♣ A♦ A♥	A♠ K♦
A♣ A♦ A♠	A♥ K♦
A♣ A♠ A♥	A♦ K♦
A♠ A♦ A♥	A♣ K♦

Is the other argument correct? How do I avoid fallacious reasoning?



REVERSIBILITY CHECK:

For each object can I reverse engineer the unique sequence of choices that constructed it?




Scheme I

1. Choose 3 of 4 aces
2. Choose 2 of the remaining cards

A♣ A♦ A♥ A♠ K♦

For this hand – you can't reverse to a unique choice sequence.

A♣ A♦ A♥	A♠ K♦
A♣ A♦ A♠	A♥ K♦
A♣ A♠ A♥	A♦ K♦
A♠ A♦ A♥	A♣ K♦





Scheme II


1. Choose 3 out of 4 aces
2. Choose 2 out of 48 non-ace cards

A♣ A♦ Q♦ A♠ K♦

REVERSE TEST: Aces came from choices in (1)
and others came from choices in (2)



Counting Poker Hands



52 Card Deck, 5 card hands

4 possible suits:

♥ ♦ ♣ ♠

13 possible ranks:

2,3,4,5,6,7,8,9,10,J,Q,K,A

Pair: set of two cards of the same rank

Straight: 5 cards of consecutive rank

Flush: set of 5 cards with the same suit

Ranked Poker Hands

Straight Flush: a straight and a flush

4 of a kind: 4 cards of the same rank

Full House: 3 of one kind and 2 of another

Flush: a flush, but not a straight

Straight: a straight, but not a flush

3 of a kind: 3 of the same rank, but not a full house or 4 of a kind

2 Pair: 2 pairs, but not 4 of a kind or a full house

A Pair

Straight Flush

9 choices for rank of lowest card at the start of the straight

4 possible suits for the flush

$$9 \times 4 = 36$$

$$\frac{36}{\binom{52}{5}} = \frac{36}{2,598,960} = 1 \text{ in } 72,193.333\dots$$

4 of a Kind

13 choices of rank

48 choices for remaining card

$$13 \times 48 = 624$$

$$\frac{624}{\binom{52}{5}} = \frac{624}{2,598,960} = 1 \text{ in } 4,165$$

Flush

4 choices of suit

$\binom{13}{5}$ choices of cards

$$4 \times 1287 = 5148$$

“but not a straight flush...”

- 36 straight flushes

5112 flushes

$$\frac{5,112}{\binom{52}{5}} = 1 \text{ in } 508.4\dots$$

Straight

9 choices of lowest card

4⁵ choices of suits for 5 cards

$$\left. \begin{array}{l} 9 \text{ choices of lowest card} \\ 4^5 \text{ choices of suits for 5 cards} \end{array} \right\} \begin{array}{l} 9 \times 1024 \\ = 9216 \end{array}$$


“but not a straight flush...”

- 36 straight flushes

9180 flushes

$$\frac{9,180}{\binom{52}{5}} = 1 \text{ in } 208.1\dots$$

Hand	Number
Straight Flush:	36
Four of a Kind:	624
Full House:	3,744
Flush:	5,112
Straight:	9,180
Three of a Kind:	54,912
Two Pair:	123,552
One Pair:	1,098,240
Nothing:	1,302,540
	2,598,960

Partition and Difference Methods**Correspondence Principle**

If two finite sets can be placed into
1-1 onto correspondence, then they
have the same size

Choice Tree**Product Rule**

Two conditions

Reverse Test

**Here's What
You Need to
Know...**

Binomial coefficient

Counting Poker Hands