



Great Theoretical Ideas In Computer Science		
Steven Rudich	CS 15-251	Spring 2005
Lecture 11	Feb 15, 2005	Carnegie Mellon University

The Mathematics Of 1950's Dating: Who wins the battle of the sexes?




Dating Scenario

- There are n boys and n girls
- Each girl has her own ranked preference list of all the boys
- Each boy has his own ranked preference list of the girls
- The lists have no ties

Question: How do we pair them off?
What criteria come to mind?


Steven Rudich: www.discretemath.com
www.rudich.net



There is more than one notion of what constitutes a "good" pairing.

- Maximizing total satisfaction
 - Hong Kong and to an extent the United States
- Maximizing the minimum satisfaction
 - Western Europe
- Minimizing the maximum difference in mate ranks
 - Sweden
- Maximizing the number of people who get their first choice
 - Barbie and Ken Land

Steven Rudich: www.discretemath.com
www.rudich.net



We will ignore the
issue of what is
“equitable”!

Steven Rudich: www.discretemath.com
www.rudich.net


Rogue Couples

Suppose we pair off all the boys and girls. Now suppose that some boy and some girl prefer each other to the people to whom they are paired. They will be called a rogue couple.



Steven Rudich: www.discretemath.com
www.rudich.net


Why be with them when we
can be with each other?



Steven Rudich: www.discretemath.com
www.rudich.net

Stable Pairings

A pairing of boys and girls is called stable if it contains no rogue couples.



Steven Rudich: www.discretemath.com
www.rudich.net

What use is fairness,
if it is not stable?

Any list of criteria for a good pairing
must include **stability**. (A pairing is
doomed if it contains a rogue couple.)

Any reasonable list of criteria must
contain the stability criterion.

Steven Rudich: www.discretemath.com
www.rudich.net

Steven's social and political wisdom:

Sustainability is a
prerequisite of *fair*
policy.

Steven Rudich: www.discretemath.com
www.rudich.net

The study of stability will be the subject of the entire lecture.

We will:


- Analyze various mathematical properties of an algorithm that looks a lot like 1950's dating
- Discover the **naked mathematical truth** about which sex has the romantic edge
- Learn how the world's largest, most successful dating service operates

Steven Rudich: www.discretemath.com
www.rudich.net

Given a set of preference lists, how do we find a stable pairing?

Steven Rudich: www.discretemath.com
www.rudich.net

Given a set of preference lists, how do we find a stable pairing?



Wait! We don't even know that such a pairing always exists!

Steven Rudich: www.discretemath.com
www.rudich.net


Given a set of preference lists, how do we find a stable pairing?



How could we change the question we are asking?

Steven Rudich: www.discretemath.com
www.rudich.net


Better Questions:



Does every set of preference lists have a stable pairing? Is there a fast algorithm that, given any set of input lists, will output a stable pairing, if one exists for those lists?

Steven Rudich: www.discretemath.com
www.rudich.net


Think about this question:



Does every set of preference lists have a stable pairing?

Steven Rudich: www.discretemath.com
www.rudich.net


Idea: Allow the pairs to keep breaking up and reforming until they become stable.




Can you argue that the couples will not continue breaking up and reforming forever?




An Instructive Variant:
Bisexual Dating




An Instructive Variant:
Bisexual Dating




An Instructive Variant:
Bisexual Dating



An Instructive Variant:
Bisexual Dating





Insight

Any proof that
heterosexual couples do not
break up and reform forever
must contain a step that
fails in the bisexual case

Steven Rudich: www.discretemath.com
www.rudich.net

Insight

If you have a proof idea that
works equally well in the hetero
and bisexual versions, then your
idea is not adequate to show
the couples eventually stop.


Steven Rudich: www.discretemath.com
www.rudich.net

The Traditional Marriage Algorithm



Steven Rudich: www.discretemath.com
www.rudich.net

The Traditional Marriage Algorithm



Traditional Marriage Algorithm

For each day that some boy gets a "No" do:

- **Morning**
 - Each girl stands on her balcony
 - Each boy proposes under the balcony of the best girl whom he has not yet crossed off
- **Afternoon (for those girls with at least one suitor)**
 - To today's best suitor: **"Maybe, come back tomorrow"**
 - To any others: **"No, I will never marry you"**
- **Evening**
 - Any rejected boy crosses the girl off his list


Each girl marries the boy to whom she just said "maybe"

Steven Rudich: www.discretemath.com
www.rudich.net

Does the Traditional Marriage Algorithm always produce a stable pairing?

Steven Rudich: www.discretemath.com
www.rudich.net

Does the Traditional Marriage Algorithm always produce a stable pairing?



Steven Rudich: www.discretemath.com
www.rudich.net

Does TMA always terminate?

- It might encounter a situation where algorithm does not specify what to do next (core dump error)
- It might keep on going for an infinite number of days

Steven Rudich: www.discretemath.com
www.rudich.net

Traditional Marriage Algorithm

For each day that some boy gets a "No" do:

- **Morning**
 - Each girl stands on her balcony
 - Each boy proposes under the balcony of the best girl whom he has not yet crossed off
- **Afternoon (for those girls with at least one suitor)**
 - To today's best suitor: "**Maybe, come back tomorrow**"
 - To any others: "**No, I will never marry you**"
- **Evening**
 - Any rejected boy crosses the girl off his list

Each girl marries the boy to whom she just said "maybe"

Steven Rudich: www.discretemath.com
www.rudich.net

Improvement Lemma: If a girl has a boy on a string, then she will always have someone at least as good on a string, (or for a husband).

- She would only let go of him in order to "maybe" someone better
- She would only let go of that guy for someone even better
- She would only let go of that guy for someone even better
- AND SO ON

Informal Induction



Improvement Lemma: If a girl has a boy on a string, then she will always have someone at least as good on a string, (or for a husband).


PROOF: Let q be the day she first gets b on a string. If the lemma is false, there must be a smallest k such that the girl has some b^* inferior to b on day $q+k$.

One day earlier, she has someone as good as b . Hence, a better suitor than b^* returns the next day. She will choose the better suitor contradicting the assumption that her prospects went below b on day $q+k$.

Formal Induction



Corollary: Each girl will marry her absolute favorite of the boys who visit her during the TMA



Steven Rudich: www.discretemath.com
www.rudich.net

Lemma: No boy can be rejected by all the girls

Proof by contradiction.

Suppose boy b is rejected by all the girls.
At that point:

- Each girl must have a suitor other than b (By Improvement Lemma, once a girl has a suitor she will always have at least one)
- The n girls have n suitors, b not among them. Thus, there are at least $n+1$ boys

Contradiction

Steven Rudich: www.discretemath.com
www.rudich.net

Theorem: The TMA always terminates in at most n^2 days

- A "master list" of all n of the boys lists starts with a total of $n \times n = n^2$ girls on it.
- Each day that at least one boy gets a "No", at least one girl gets crossed off the master list
- Therefore, the number of days is bounded by the original size of the master list. In fact, since each list never drops below 1, the number of days is bounded by $n(n-1) = n^2$.


Steven Rudich: www.discretemath.com
www.rudich.net

Great! We know that TMA will terminate and produce a pairing.

But is it stable?


Steven Rudich: www.discretemath.com
www.rudich.net

Theorem: Let T be the pairing produced by TMA. T is stable.



Steven Rudich: www.discretemath.com
www.rudich.net

Theorem: Let T be the pairing produced by TMA. T is stable.



Steven Rudich: www.discretemath.com
www.rudich.net

Theorem: Let T be the pairing produced by TMA. T is stable.

- Let b and g be any couple in T .
- Suppose b prefers g^* to g . We will argue that g^* prefers her husband to b .
- During TMA, b proposed to g^* before he proposed to g . Hence, at some point g^* rejected b for someone she preferred. By the Improvement lemma, the person she married was also preferable to b .
- Thus, every boy will be rejected by any girl he prefers to his wife. T is stable.

Steven Rudich: www.discretemath.com
www.rudich.net

Opinion Poll



Steven Rudich: www.discretemath.com
www.rudich.net

Forget TMA for a moment

How should we define what we mean when we say "the optimal girl for boy b "?

Flawed Attempt:
"The girl at the top of b 's list"

Steven Rudich: www.discretemath.com
www.rudich.net

The Optimal Girl

A boy's optimal girl is the highest ranked girl for whom there is some stable pairing in which the boy gets her.

She is the best girl he can conceivably get in a stable world. Presumably, she might be better than the girl he gets in the stable pairing output by TMA.

Steven Rudich: www.discretemath.com
www.rudich.net

The Pessimal Girl

A boy's pessimal girl is the lowest ranked girl for whom there is some stable pairing in which the boy gets her.

She is the worst girl he can conceivably get in a stable world.

Steven Rudich: www.discretemath.com
www.rudich.net

Dating Heaven and Hell

A pairing is male-optimal if every boy gets his optimal mate. This is the best of all possible stable worlds for every boy simultaneously.

A pairing is male-pessimal if every boy gets his pessimal mate. This is the worst of all possible stable worlds for every boy simultaneously.

Steven Rudich: www.discretemath.com
www.rudich.net

Dating Heaven and Hell

A pairing is male-optimal if every boy gets his optimal mate. Thus, the pairing is simultaneously giving each boy his optimal.

Is a male-optimal pairing always stable?

Steven Rudich: www.discretemath.com
www.rudich.net

Dating Heaven and Hell

A pairing is female-optimal if every girl gets her optimal mate. This is the best of all possible stable worlds for every girl simultaneously.

A pairing is female-pessimal if every girl gets her pessimal mate. This is the worst of all possible stable worlds for every girl simultaneously.

Steven Rudich: www.discretemath.com
www.rudich.net

The Naked Mathematical Truth!

The Traditional Marriage Algorithm always produces a male-optimal, female-pessimal pairing.

Steven Rudich: www.discretemath.com
www.rudich.net

Theorem: TMA produces a male-optimal pairing

- Suppose, for a contradiction, that some boy gets rejected by his optimal girl during TMA. Let t be the earliest time at which this happened.
- In particular, at time t , some boy b got rejected by his optimal girl g because she said "maybe" to a preferred b^* . By the definition of t , b^* had not yet been rejected by his optimal girl.
- Therefore, b^* likes g at least as much as his optimal.

Steven Rudich: www.discretemath.com
www.rudich.net

Some boy b got rejected by his optimal girl g because she said "maybe" to a preferred b^* . b^* likes g at least as much as his optimal girl.

There must exist a stable pairing S in which b and g are married.

- b^* wants g more than his wife in S
 - g is as at least as good as his best and he does not have her in stable pairing S
- g wants b^* more than her husband in S
 - b is her husband in S and she rejects him for b^* in TMA

Steven Rudich: www.discretemath.com
www.rudich.net

Some boy b got rejected by his optimal girl g because she said "maybe" to a preferred b^* . b^* likes g at least as much as his optimal girl.

There must exist a stable pairing S in which b and g are married.


- b^* wants g more than his wife in S
 - g is as at least as good as his best and he does not have her in stable pairing S
- g wants b^* more than her husband in S
 - b is her husband in S and she rejects him for b^* in TMA

Contradiction of the stability of S .

What proof technique did we just use?

Steven Rudich: www.discretemath.com
www.rudich.net

What proof technique did we just use?



Theorem: The TMA pairing, T , is female-pessimal.

We know it is male-optimal. Suppose there is a stable pairing S where some girl g does worse than in T .

Let b be her mate in T .

Let b^* be her mate in S .

- By assumption, g likes b better than her mate in S
- b likes g better than his mate in S
 - We already know that g is his optimal girl
- **Therefore, S is not stable.**

Contradiction

Steven Rudich: www.discretemath.com
www.rudich.net

Advice to females

Learn to make the first move.

Steven Rudich: www.discretemath.com
www.rudich.net

The largest, most successful
dating service in the world
uses a computer to run TMA !

Steven Rudich: www.discretemath.com
www.rudich.net

REFERENCES

D. Gale and L. S. Shapley, *College admissions and the stability of marriage*, American Mathematical Monthly 69 (1962), 9-15

Dan Gusfield and Robert W. Irving, *The Stable Marriage Problem: Structures and Algorithms*, MIT Press, 1989

Steven Rudich: www.discretemath.com
www.rudich.net