

Cost-Sharing Mechanisms for Network Design*

Anupam Gupta[†]

Aravind Srinivasan[‡]

Éva Tardos[§]

Abstract

We consider a single-source network design problem from a game-theoretic perspective. Gupta, Kumar and Roughgarden (*Proc. 35th Annual ACM STOC*, pages 365–372, 2003) developed a simple method for a single-source rent-or-buy problem that also yields the best-known approximation ratio for the problem. We show how to use a variant of this method to develop an approximately budget-balanced and group strategyproof cost-sharing method for the problem.

The novelty of our approach stems from our obtaining the cost-sharing methods for the rent-or-buy problem by carefully combining cost-shares for the simpler Steiner tree problem. Our algorithm is conceptually simpler than the previous such cost-sharing method due to Pál and Tardos (*Proc. 44th Annual FOCS*, pages 584–593, 2003), and improves the previously-known approximation factor of 15 to 4.6.

1 Introduction

This paper studies the problem of giving good *cost-sharing mechanisms* for a single-source network design problem. Imagine a general network design problem, where the participants (or *agents*) want to build a network connecting them to a common source (a server); however, they are autonomous and behave in a selfish (but non-malicious) fashion. Informally, a cost-sharing mechanism builds a network, and allocates the cost incurred among the agents, so that no group of agents is charged too much, thus precluding the possibility of their being unhappy and trying to secede from the system.

The type of problem we consider is where we are given an undirected graph $G = (V, E)$, and a set of *demands* $D \subseteq V$ that want to connect to a common source r . We want to devise an algorithm that builds a cheap network on the demands D , and also specify what portion of its cost is paid by which of the participants in the network; moreover, this should be done in a manner that ensures that the cost paid by any subset of the participants is “fair”.

To make all this precise, let us consider a network design game on a graph $G = (V, E)$. The cost-shares will be specified by a function $\xi : 2^V \times V \rightarrow \mathbb{R}_{\geq 0}$, where $\xi(D, i)$ will specify the cost share paid by the player $i \in V$ when the actual set of demands is $D \subseteq V$. To ensure that players are not penalized when they do not participate

*A preliminary version of this work appears in the *Proc. International Workshop on Approximation Algorithms for Combinatorial Optimization Problems*, 2004. This research was done in part during the IMA Workshop on *Network Management and Design* at the University of Minnesota, April 2003.

[†]Department of Computer Science, Carnegie Mellon University, Pittsburgh, PA 15232. Email: anupamg@cs.cmu.edu.

[‡]Department of Computer Science and University of Maryland Institute for Advanced Computer Studies, University of Maryland at College Park, College Park, MD 20742. This material is based upon work supported in part by the National Science Foundation under Grant No. 0208005 and ITR Award CNS-0426683. srin@cs.umd.edu

[§]Department of Computer Science, Cornell University, Ithaca, NY 14853. eva@cs.cornell.edu. Research supported in part by ONR grant N00014-98-1-0589, and NSF grants CCR-0311333 and CCR-0325453.

in the game, we require that $\xi(D, i) = 0$ for $i \notin D$. For a particular set D of demands, we use $\text{OPT}(D)$ to denote the minimum cost network serving the users in the set D , and use $\text{Alg}(D)$ to denote the cost (or expected cost, in the case of randomized algorithms) of the network computed by our algorithm. We will be concerned with designing algorithms and cost-sharing functions ξ that satisfy the following properties:

1. (**β -approximate budget-balance**) For any set of demands D , we require that

$$\text{OPT}(D) \geq \sum_{i \in D} \xi(D, i) \geq \text{Alg}(D)/\beta$$

for some given parameter $\beta \geq 1$. Equivalently, by multiplying the shares by β , we could require that the cost-shares are at least the total cost of the solution found by the algorithm, but do not exceed $\beta \text{OPT}(D)$. If $\beta = 1$, we say that the cost-sharing function ξ is *budget-balanced*.

2. (**cross-monotonicity**) For any $A \subseteq D$ and any $i \in A$, $\xi(D, i) \leq \xi(A, i)$. I.e., the cost-share of any fixed demand point i should not increase due to other demands entering the system. This property is also known under the name of population-monotone.

It is well-known that the two properties above immediately imply the following desirable property (see, e.g., [8] or Lemma 3.1, for a proof):

3. (**fairness**) For any $A \subseteq D$, $\sum_{i \in A} \xi(D, i) \leq \text{OPT}(A)$. I.e., the cost paid by any subset of people should not exceed the optimal cost of connecting them alone and hence they have no incentive to secede. This property is often referred to as *competitiveness*.

Cross-monotonicity is a key ingredient used in solving the following type of mechanism design problems/games: consider the network design problem with a set of demand nodes D , with each user (or demand) d having an associated utility u_d . Since the users have limited utilities, the service provider has to now decide which subset of customers it must serve, in addition to designing the network and deciding how to share the cost between the served customers. A mechanism for solving this problem is called *group strategyproof* if no subset of users has an incentive to deviate from the protocol (e.g., by misreporting their utility) in the hope of improving the outcome for themselves (e.g., receiving the service at a cheaper cost). Moulin and Shenker [8] show that having a cross-monotone cost-sharing method for a problem naturally gives rise to a group strategyproof mechanism for the problem in the following way. We start with all the customers; if there is some customer whose cost share (computed w.r.t. the current set of customers) exceeds its utility, we drop it from the set, recompute cost shares and repeat. At the end, we are left with the desired set of customers and their cost-shares.

As an example of cross-monotonicity, let us consider the MINIMUM SPANNING TREE game on the complete graph $G = (V, E)$ with edge weights, given by a set of players D and a *root* $r \notin D$; the objective is to find the cheapest tree $\text{MST}(D)$ spanning D and r . It must be emphasized that this game does not allow the use of Steiner vertices in $\text{MST}(D)$, and r is not a player, and hence should have no cost-share. It is not difficult to verify that a budget-balanced and fair cost-sharing scheme for this game can be found thus: find an MST, root it at r , and set the cost-share for vertex $i \in D$ to be the cost of the edge from i to its parent. However, this scheme is not cross-monotone, and getting a cross-monotone cost-sharing scheme for the minimum spanning tree problem takes more work. Kent and Skorin-Kapov [6] and Jain and Vazirani [9] developed such budget-balanced and cross-monotone cost-sharing schemes for the spanning tree game using a directed branching game; let us denote this cost-sharing scheme by ξ_{MST} . It is well-known that a min-cost spanning tree is a 2-approximation to the min-cost Steiner tree; hence once can infer that the values $\frac{1}{2}\xi_{\text{MST}}$ serve also as cross-monotone, 2-approximately budget-balanced cost-shares for the corresponding Steiner tree game.

The Single-Source Rent-or-Buy Game In this paper we will consider the *Single-Source Rent-or-Buy* Network Design game; this combines features of Steiner tree and shortest paths. The game is defined as follows: we are given a complete undirected graph $G = (V, E)$ with edge costs c_e satisfying the triangle inequality,¹ a special *source* (or *root*) vertex r , and an integer parameter $M \geq 1$. There are also a set of *users* (also called *players* or *demands*) $D \subseteq V$, each of which is identified with a vertex of the graph. We assume that there is a unique player at each vertex. Our results can be extended (with a slight loss in the performance guarantee) to multiple players at each vertex, each with a *weight* d_j indicating its amount of demand; we sketch how to do this in Section 6.2.

The objective is to connect each player j to the source r via some path P_j , on which one unit of bandwidth has been allocated. What makes the game interesting is that aggregating paths is beneficial, in tune with the idea of economies of scale. Hence there are two different actions that can be performed by each edge: either the edge can be *bought* at cost Mc_e , but then an arbitrary amount of bandwidth can be sent on that edge (and hence an arbitrary number of paths P_j can use it); or bandwidth on the edge can be *rented* at cost c_e per unit bandwidth (and hence if the paths for some set S of players were using the edge, then the edge would cost $c_e \times |S|$; in the general weighted case, the cost would be c_e times the total weight of the players in S). As usual, any edge can be used for any of the users. Our main theorem is the following:

Theorem 1.1 *There is a cross-monotone cost-sharing method for the Single-Source Rent-or-Buy network design game that is also β -budget-balanced, where $\beta = 4.6$. Furthermore, these cost-shares can be computed in deterministic polynomial time.*

This improves on the results of Pál and Tardos [13], who gave a 15-approximate cross-monotone cost-sharing scheme for the problem. They use a primal-dual algorithm to build the network and obtain cost-shares.

We construct the claimed cost-shares based on expected costs in the approximation algorithm of Gupta et al. [5]. Loosely speaking, that algorithm works by randomly reducing the network design problem to computing Steiner trees over subsets of terminals; we show that if we use the function $\frac{1}{2}\xi_{MST}$ to allocate approximate cost-shares to each vertex in this resulting Steiner game, then the expected cost share of a vertex (taken over the random reduction) gives us the cross-monotone cost-shares claimed in Theorem 1.1. However, since the variances of the random variables involved may be large in general, computing these expected costs in polynomial time is non-trivial. (In fact, a similar method to obtain cost-shares was proposed independently by Leonardi and Schäfer [11], who showed a budget-balance factor of $4(1 + \epsilon)$ for any $\epsilon > 0$ for their cost-shares; however, they left open the problem of the efficient computation of these costs.) To do this computation, we have to give a new alternative analysis of the algorithm of [5]; this analysis allows us to give a derandomization using a small sample space, which has some additional useful properties that help ensure cross-monotonicity.

As is standard, we often use the symbol e to denote an edge in G . The same symbol will also be used in some places to denote the base of the natural logarithm; since the two usages are sufficiently distinct, we hope this does not cause confusion. Logarithms will be to the base two, unless specified otherwise. The rest of this paper is organized as follows. The basic algorithm and the idea for computing cost-shares are discussed in Section 2; Section 3 proves some basic properties of the cost-sharing scheme, such as cross-monotonicity. Section 4 gives a way of analyzing our algorithm—bounding β , in particular—that is different from the one of [5]. This proof is used as a starting-point in Section 5 to show that our estimate of β from Section 4 changes negligibly in our derandomized algorithm. Finally, further extensions are described in Section 6.

¹Since we can work with the metric completion of the input graph instead, the assumption of the triangle inequality is without loss of generality.

2 The Algorithm and the Cost-Sharing

We will use an algorithm `SimpleCFL` suggested by Gupta et al. [5] for the equivalent problem of connected facility location (without facility costs). First, let us recall the algorithm from [5]. Here $\alpha > 0$ is a constant that will be chosen later.

- S1. *Mark* each demand j independently with probability α/M , and let D' be the set of marked demands.
- S2. Construct a minimum spanning tree T on $F = D' \cup \{r\}$. (This is a 2-approximate Steiner tree on F .) The elements of F are called the *open facilities*.
- S3. Assign each demand $j \in D$ to its closest open facility $i(j)$ in F .

The algorithm suggests a simple and intuitive idea for the cost-shares: each player pays a cost proportional to the *expected* cost incurred by it on running the above algorithm. For a given a sequence of coin-tosses, the players in D' will pay for *buying* the MST, where their shares are derived from the cross-monotone cost-sharing scheme ξ_{MST} for the MST problem given in [6, 9]. All other players (players in $D \setminus D'$) will pay for *renting* their shortest paths to F . The cost-shares defined for a particular set of coin-tosses exactly cover the cost of the solution built. From [5] we know that this algorithm is a β -approximation algorithm for some constant β . To get β -approximate cost-shares we divide the above defined shares by β . Formally:

$$\xi(D, j) = \frac{1}{\beta} \mathbb{E}[M \xi_{MST}(F, j) + \ell(j, F)], \quad (2.1)$$

where $\ell(j, S)$ is the length of a shortest-path from j to the closest vertex of the set S , and the expectation is over the coin tosses. (Note that the set $F = D' \cup \{r\}$ depends on the coin tosses.)

Outline of the Proof There are two parts to proving the result: we first have to show that the properties of the cost-shares claimed in Theorem 1.1 are indeed satisfied; i.e., they are β -budget-balanced and cross-monotone. These properties are proved in Section 3.

The technically involved part of the proof involves showing that our cost-shares can be computed in (deterministic) polynomial time. A little thought will convince the reader that this is not obvious, even with the results of Gupta et al. [5]. Indeed, we need to estimate the expectations in (2.1) for each of the players, but the random variables in (2.1) do not have small variance in general. Furthermore, it is unclear how to derandomize the proof of [5], since it relies on some severe conditioning. To take care of this problem, we give a proof of the performance guarantee of the `SimpleCFL` algorithm different from the one given by [5]; this is done in Section 4. Our new proof will yield a somewhat worse constant than that of [5], but allow us to derandomize the algorithm by marking the demands in Step (S1) in t -wise independent fashion for a constant t ; this appears in Section 5. Using t -wise independent random choices in Step (S1) allows us to use a polynomial-sized sample space, letting us compute expectations such as those of (2.1) in (deterministic) polynomial time, by considering all the points in the sample space. However, we need to make sure that the properties of the expectation proved in Section 3 (fairness and cross-monotonicity) also hold for the expectation using t -wise independent random choices. Interestingly, the properties of a *particular* construction of t -wise independent random variables turns out to be crucial, as described in the proof of Theorem 3.5.

3 Properties of the Cost-Sharing Scheme

Recall that given an instance of the game specified by a set of demands D , a *cost-sharing scheme* for the game is simply a function $\xi(D, i)$ (with $\xi(D, i) = 0$ for $i \notin D$). We now need to show that the function defined in (2.1) is a cost-sharing scheme with the properties we care about, namely, approximate budget-balance, fairness, and cross-monotonicity. As mentioned in the introduction, it is well-known that these three properties are not independent, and that a cross-monotone and approximately budget balanced cost-sharing is always fair.

Lemma 3.1 *If a cost-sharing function ξ is approximately budget balanced and cross-monotone then it is also fair.*

Proof: Consider a subset $A \subseteq D$. By cross-monotonicity, it follows that $\sum_{i \in A} \xi(D, i) \leq \sum_{i \in A} \xi(A, i)$. Now using the upper bound in the approximate budget-balance property gives us $\sum_{i \in A} \xi(A, i) \leq \text{OPT}(A)$, and completes the proof of the lemma. ■

We will need the following facts about the cost-sharing scheme ξ_{MST} :

Theorem 3.2 ([6, 9]) *There exists an efficiently computable cost-sharing scheme ξ_{MST} for the Minimum Spanning Tree game that is budget-balanced and cross-monotone.*

Since the spanning tree connects D and r , we will use the notation $\xi_{MST}(D \cup \{r\}, i)$ and $\xi_{MST}(D, i)$ interchangeably; however, the results of [6, 9] will always ensure that $\xi_{MST}(D \cup \{r\}, r) = 0$.

To prove approximate budget-balance, we will need the following result bounding the performance of the algorithm SimpleCFL; its proof appears in Sections 4 and 5.

Theorem 3.3 *The expected cost of the algorithm SimpleCFL on a set of demands D is at most $\beta \text{OPT}(D)$ (where $\beta = 4.6$), even if the demands are marked in a t -wise independent fashion in Step (S1), for a suitably large constant t .*

Note that this is a stronger form of the performance guarantee proved in [5] (albeit with a worse constant), since it requires only constant-wise independence of the random variables. Armed with this result, we are now in a position to prove the properties of the function ξ defined in (2.1), and thus to prove Theorem 1.1.

Notation. In our randomized algorithm, Y_j will denote the indicator random variable for demand j getting marked. Given a sample point ω (i.e., the underlying sequence of coin-tosses) in our probability space, $Y_j(\omega)$ will be the following deterministic quantity: 1, if j gets marked due to this sequence of coin-tosses, and 0 otherwise.

The above two theorems imply that ξ is β -approximately budget-balanced – for any random sample ω , ξ shares exactly the cost of the solution built:

Lemma 3.4 *The function ξ is β -approximately budget-balanced.*

Proof: Consider the cost-shares $\beta \xi$. For a fixed random sample ω , let F_ω denote the set of facilities; i.e., $F_\omega = \{r\} \cup \{j \in D \mid Y_j(\omega) = 1\}$. Also let $c(\text{MST}(F_\omega))$ denote the cost of an MST for F_ω . From our definition of $\xi(D, i)$, we get the following:

$$\sum_{i \in D} \beta \xi(D, i) = \mathbb{E}_\omega \left[M \sum_{i \in D} \xi_{MST}(F_\omega, i) + \sum_{i \in D} \ell(i, F_\omega) \right] = \mathbb{E}_\omega \left[M \times c(\text{MST}(F_\omega)) + \sum_{i \in D} \ell(i, F_\omega) \right], \quad (3.2)$$

using the facts that ξ_{MST} is a cost-share function and hence $\xi_{MST}(F_\omega, i) = 0$ for $i \notin F_\omega$, and that ξ_{MST} is budget-balanced and thus $\sum_{i \in F_\omega} \xi_{MST}(F_\omega, i) = c(\text{MST}(F_\omega))$.

Now it is easy to see that the cost-shares $\beta \xi$ recover the cost of the solution constructed, and hence are at least $\text{Alg}(D)$. This is true, as for each element of the sample space ω , $M \times c(\text{MST}(F_\omega)) + \sum_i \ell(i, F_\omega)$ is the cost of the solution which opens facilities at F_ω .

To see that $\sum_{i \in D} \xi(D, i) \leq \text{OPT}(D)$, note that the final expression in (3.2) is the expected cost of running **SimpleCFL** on the set of demands D , and hence Theorem 3.3 implies that the expectation is no more than $\beta \text{OPT}(D)$, and thus $\sum_{i \in D} \xi(D, i) \leq \text{OPT}(D)$. \blacksquare

3.1 Details of the limited-independence marking

Before presenting the specific type of t -wise independent random marking that we will use, it would be useful to see how β depends on α ; recall that α was a parameter used to determine the marking probability in Step (S1) of the algorithm **SimpleCFL**. We show first in bound (4.17) of Section 4.2 that

$$\beta \leq \max \left\{ 2(1 + \alpha), \frac{2(2e^{2\alpha} + e^\alpha - 2)}{e^{2\alpha} - 1} \right\} \quad (3.3)$$

under fully-independent marking. We then show in Section 5.2 that this bound changes negligibly if t is a sufficiently large constant: if $t = a \log(1/\epsilon)$ where a is an absolute constant, then the r.h.s. of (3.3) gets multiplied at most by $(1 + \epsilon)$. Thus, setting $\alpha = 1.296$, we will be able to guarantee that $\beta \leq 4.6$.

In order to prove that the expectation using t -wise independent random marking is cross-monotone, we need a particular choice of random marking. Let \mathbb{F} be a field of size $\geq n$, with $|\mathbb{F}|$ chosen large enough so that $\lceil \alpha |\mathbb{F}| / M \rceil / |\mathbb{F}|$ is sufficiently close to α / M ; in particular, suppose $|\mathbb{F}| \geq \Omega(M \log n)$, say. Let the elements of the field be $\{a_1, a_2, \dots, a_{|\mathbb{F}|}\}$, and let the vertices V of the graph be labeled by the first n elements $\{a_1, a_2, \dots, a_n\}$. Let \mathcal{S} be any pre-specified subset of \mathbb{F} with $|\mathcal{S}| = \lceil \alpha |\mathbb{F}| / M \rceil$. To get a sample, generate $\omega = (x_0, x_1, \dots, x_{t-1})$ uniformly at random from \mathbb{F}^t , and define Y_i to be 1 if $\sum_{j=0}^{t-1} x_j a_i^j$ lies in \mathcal{S} , and $Y_i = 0$ otherwise. By construction, we have $\Pr [Y_i = 1] \sim \alpha / M$. In particular, since $|\mathbb{F}| \geq \Omega(M \log n)$, we have

$$\alpha / M \leq \Pr [Y_i = 1] \leq \alpha / M + 1 / |\mathbb{F}| = (\alpha + o(1)) / M;$$

by (3.3) and the few sentences following it, we see that this replacement of α by $\alpha + o(1)$ impacts β only by an additive $o(1)$ factor. Therefore, in all our arguments below, we will assume that $\Pr [Y_i = 1] = \alpha / M$ for all $i \in D$. It is well-known that the Y_i generated as above are all t -wise independent; see, e.g., [1, 7]. Note that the above distribution is generating n coin tosses Y_i , while we need only $|D|$ of them; however, we can just ignore all the Y_j for $j \notin D$.

For a fixed $\omega \in \mathbb{F}^t$, let F_ω denote the set of facilities; i.e., $F_\omega = \{r\} \cup \{j \in D \mid Y_j(\omega) = 1\}$.

Theorem 3.5 *Assuming that the random marking is done using the t -wise independent random variables explained above, or using independent random variables, then the function ξ is cross-monotone, i.e., $\xi(A, i) \geq \xi(D, i)$ for any $A \subseteq D$.*

Proof: The particular type of t -wise independent distribution that we use has the following natural but crucial property: for any choice of $\omega \in \mathbb{F}^t$, if the set of marked demands in the run of **SimpleCFL** on A is $A'(\omega)$ and the set of marked demands in the run on D is $D'(\omega)$, then $A'(\omega) \subseteq D'(\omega)$. (To our knowledge, this crucial property need not necessarily hold for all t -wise independent distributions.) Define the joint probability $p(F, E)$ to be the probability of selecting an element of the sample space ω such that the marked demands in the run of **SimpleCFL** on A is F and on D it is E . Note that $p(F, E) > 0$ only if $F \subseteq E$. A joint probability with this property can also be defined for the case when the marking is fully independent.

With this additional property, we can prove that the scheme ξ is cross-monotone. To see this, note that

$$\xi(A, i) = \frac{1}{\beta} \sum_{F, E} p(F, E) [M \xi_{MST}(F, i) + \ell(i, F)] \geq \frac{1}{\beta} \sum_{F, E} p(F, E) [M \xi_{MST}(E, i) + \ell(i, E)] = \xi(D, i),$$

where the inequality uses the fact that the support of $p(F, E)$ is only on pairs with $F \subseteq E$, and the cross-monotonicity of ξ_{MST} . \blacksquare

4 A New Analysis of the Performance of SimpleCFL

Suppose first that the marking is done fully independently in Step (S1); we now give a way of analyzing SimpleCFL that is different from the one of [5]. We will then use this proof in Section 5 to show that our estimate of β changes negligibly when we conduct a t -wise independent marking for a large-enough (but constant) value of t . Recall that a solution to the Connected Facility Location Problem with demands D is given by the facilities F and the Steiner tree on them, with the cost being $\sum_{j \in D} \ell(j, i^*(j)) + M c(T)$. Of the two terms in the sum, the former cost is referred to as the *connection* cost, and the latter is the *Steiner* cost. Let OPT be an optimal solution with facilities F^* and tree T^* , and let C^* and $S^* = M c(T^*)$ be the connection and Steiner costs in OPT; also let $Z^* = C^* + S^*$.

The bound for the Steiner cost S is the same as in [5]: we consider the optimal solution OPT, and compute the expected cost assuming that we construct the Steiner tree in Step (S2) using the paths of the optimal solution. We present the formal proof in Lemma 4.1; part (a) of this lemma will only be required in Section 5. The random variables Y_j are as defined in Section 3.1.

Lemma 4.1 (a) *The expected cost S of Step (S2) of SimpleCFL is bounded by a non-negative linear combination of the values $\{E[Y_j] : j \in D\}$; (b) this non-negative linear combination is at most*

$$2 (S^* + \alpha C^*). \quad (4.4)$$

Proof: Consider the optimal solution OPT: it connects each $j \in D$ to a facility $i^*(j) \in F^*$ via a shortest path, and hence pays a connection cost of $C^* = \sum_{j \in D} \ell(j, i^*(j))$. Let F be as defined in Step (S2). Define a graph connecting F thus: buy all the edges in T^* , and then for each $j \in F$, buy the edges in the shortest path from j to $i^*(j)$ used by OPT. Note that we pay $M c_e$ for each edge e in Step (S2), and hence pay $M c(T^*) = S^*$ for edges of T^* . For the random paths we add, the expected cost of the path between j and $i^*(j)$ is $E[Y_j] \times M \ell(j, i^*(j))$. Thus, the expected total cost of this graph that connects up F is at most

$$S^* + \sum_{j \in D} E[Y_j] \times M \ell(j, i^*(j)).$$

Since we buy a minimum spanning tree in Step (S2), the expected buying cost of our tree T is at most twice the expression just seen; i.e., at most

$$2 (S^* + \sum_{j \in D} E[Y_j] \times M \ell(j, i^*(j))), \quad (4.5)$$

proving part (a) of the Lemma. Since $E[Y_j] = \alpha/M$, we get part (b) of the lemma. \blacksquare

Massaging the optimal solution: Before we proceed to bound the connection costs, let us modify OPT to get another solution OPT', which costs more than OPT, but which has some more structure that allows us to complete the argument. The new solution has the following properties:

P1. OPT' opens facilities at $F' \subseteq F^*$, and these are connected by a cycle T' instead of a tree T^* ; each demand j is now connected to the facility $i'(j)$, which is not necessarily its closest facility.

P2. The number of demands assigned to a facility in F' , except perhaps the root r , is a multiple of M . (We will call this the *mod- M* constraint).

Lemma 4.2 *There is a solution OPT' with the properties above, which has connection cost $C' \leq C^* + S^*$ and Steiner cost $S' \leq 2S^*$. Hence, the total cost of OPT' is $Z' \leq C^* + 3S^*$.*

Proof: Modifying the assignment of demands to nodes in F^* to satisfy the mod- M constraint is fairly simple. Given the tree T^* , we process this in a two-pass fashion. The first pass goes bottom-up, starting at the leaves which must all be in F^* , and making sure that each node satisfies the mod- M property by sending some $b < M$ demand from the node to its parent in the tree. OPT has nodes that are not in F^* , and hence cannot be facilities in F' . We eliminate these nodes during a top-down pass, where we distribute their demand among their children. The details are as follows.

Given the tree T^* , we process this in a two-pass fashion. The first pass goes bottom-up, starting at the leaves which must all be in F^* . Consider a vertex all of whose descendants satisfy the mod- M property, and let x be the demand currently assigned to it. If $x = aM + b$, where $b < M$, we send b units of demand from x to its parent. The number of demands assigned to each is now a multiple of M . Furthermore, the increase in connection cost is $b < M$ times the distance between x and its parent; this ensures that every node (except possibly r) has demands that are a multiple of M .

Of course, we may now have demands gathering at nodes of OPT that are not in F^* ; handling this is fairly simple in a top-down fashion. We will send some of the demand back down the tree from parents to children; however, we maintain the invariant that at most M units of demand will be sent down an edge. Let a vertex $v \notin F^*$ have aM demand (and let us assume we have handled all its ancestors). Since at least $(a - 1)M$ of this demand must have been collected in the bottom-up pass, and each child could have only sent $< M$ demand upwards, there must be $\geq a$ children of v sending demands to it. We choose any a of these children and send back M units of demand along those edges.

To complete the argument, note that the net demand sent along an edge of T^* was either b units upwards, or $M - b$ units downwards, and hence the cost of this movement can be changed to S^* , thus obtaining that $C' \leq C^* + S^*$. Furthermore, since we finally do not use any new facilities, and can even close the facilities with no demands assigned to it, $F' \subseteq F^*$. Finally, the cycle property is obtained by taking an Eulerian tour of T^* and replacing it by a cycle by short-cutting repeated vertices using the metric property. This at most doubles the Steiner cost, giving $S' \leq 2S^*$. ■

Back to Bounding the Connection Cost: From now on, we shall only consider the modified solution OPT' . For simplicity, by making copies of nodes if necessary, we can assume that each node in F' has been assigned M demands. Recall that T' is a cycle which contains r . (See Figure 1.) Starting from r , let us name the facilities on T' in (say) clockwise order $r = f_0, f_1, \dots, f_k, f_{k+1} = r$. (Hence referring to f_l makes sense for all l , since we can just consider $l \bmod (k + 1)$.)

Let D_l be the set of demands assigned to f_l , and hence $|D_l| = M$ for $l \neq 0$ by assumption (P2). Let us abuse notation and add r to D_0 . Let P'_l be the portion of the cycle T' joining f_l and f_{l+1} , and let $c(P'_l)$ denote the length of P'_l , and hence $S' = M \sum_{l=0}^k c(P'_l)$. Let C'_l be the total connection cost of the demands in D_l in the solution OPT' (with $C' = \sum_{l=0}^k C'_l$). Our algorithm chooses, in Step (S3), the cheapest assignment of demands to nodes in F . Thus, to bound the expected connection cost, it suffices to bound the expected connection cost of an arbitrary experiment that assigns demands to nodes in F . We present one such experiment next in Section 4.1,




Figure 1: *The transformed instance OPT'*

and analyze it; in Section 4.2, we present and briefly analyze an improved experiment. We have chosen this approach since there are several details to be handled in Section 4.2; handling these seems easier if we start with the simpler approach of Section 4.1 as a warmup.

4.1 A Candidate Assignment

Here we consider the following experiment: if $D_l \cap F \neq \emptyset$, then we assign *all* the demands in D_l to the element in $D_l \cap F$ that is closest to f_l . Else we have to assign these demands to some other open facility (i.e., “go outside D_l ”). In this case, consider the smallest $t > 0$ such that $D_{l+t} \cap F \neq \emptyset$; let $s = l + t$. Note that $r \in D_0$, and so $t \leq k$. We now send all the demands in D_l to the facility in D_s closest to f_s . If we assign demands in D_l to a marked node in D_{l+t} , then a path for each demand in D_l goes through $f_l, P'_l, P'_{l+1}, \dots, P'_{l+t-1}$, and then from f_{l+t} to the element in $D_{l+t} \cap F$ closest to it. We will bound the expected cost of these paths, which in turn will bound the expected connection cost of SimpleCFL.

Let us define some random variables. Let X_i be the indicator variable that $D_i \cap F = \emptyset$ in our algorithm. (Note that $X_0 = 0$ with probability 1.) Let A_i be the distance from f_i and the closest element of $D_i \cap F$; if $D_i \cap F$ is empty, then A_i can be set to 0. By the above arguments, the assignment cost of the M demands in D_l is at most

$$C'_l + M \sum_{i=l}^k (X_l \cdots X_i) c(P'_i) + M \sum_{i=l}^k (X_l \cdots X_{i-1}) (1 - X_i) A_i \quad (4.6)$$

Indeed, the first term is the distance traveled by the demands in D_l to reach f_l , the second term expresses the fact that demands use P_i (and pay $M c(P'_i)$) if and only if D_l, \dots, D_i all do not intersect F , and the third term implies that we incur a cost of $M A_i$ if we assign demands in D_l to the closest member of $D_i \cap F$.

Note that X_i and X_j are independent for $i \neq j$, and that $E[X_i] = (1 - \alpha/M)^M = \mathbf{q}$. To bound $M \cdot E[(1 - X_i)A_i]$, let the distances from f_i to the members of D_i be

$$0 \leq a_1 \leq a_2 \leq \dots \leq a_M, \text{ with } \sum_j a_j = C'_i. \quad (4.7)$$

Now,

$$E[(1 - X_i)A_i] = \sum_{j=1}^M a_j \times \alpha/M \times (1 - \alpha/M)^{j-1}.$$

Note that the coefficients of a_j decrease as j increases, and hence subject to the constraints (4.7), the expectation is maximized when all the a_j 's are equal to C'_i/M . This yields

$$\mathbb{E}[(1 - X_i)A_i] \leq (C'_i/M) [1 - (1 - \alpha/M)^M] = (C'_i/M) (1 - \mathbf{q}). \quad (4.8)$$

Let C_l be the expected connection cost of the demands in D_l in our current experiment. Combining the above-mentioned facts with the inequality (4.6), we get that for $l > 0$,

$$C_l \leq C'_l + M \sum_{i=l}^k c(P'_i) \mathbf{q}^{i-l+1} + \sum_{i=l}^k C'_i (1 - \mathbf{q}) \mathbf{q}^{i-l}. \quad (4.9)$$

Note that $C_0 = C'_0$; adding this to the sum of (4.9) over all l , the total expected connection cost is

$$C \leq \sum_{l=0}^k C'_l + M \sum_{i=1}^k c(P'_i) \sum_{l=1}^i \mathbf{q}^{i-l+1} + \sum_{i=1}^k C'_i (1 - \mathbf{q}) \sum_{l=1}^i \mathbf{q}^{i-l} \quad (4.10)$$

$$\begin{aligned} &\leq C' + M \sum_{i=1}^k c(P'_i) \frac{\mathbf{q}}{1 - \mathbf{q}} + \sum_{i=1}^k C'_i (1 - \mathbf{q}) \frac{1}{1 - \mathbf{q}} \\ &\leq 2C' + \frac{\mathbf{q}}{1 - \mathbf{q}} S' \end{aligned} \quad (4.11)$$

However, $\mathbf{q} \leq 1/e^\alpha$, and hence the second term is at most $1/(e^\alpha - 1)$. Now using Lemma 4.2 to replace C' by $C^* + S^*$ and S' by $2S^*$, we see that the expected connection cost of our algorithm can be bounded by:

$$C \leq 2C^* + \frac{2e^\alpha}{e^\alpha - 1} S^*. \quad (4.12)$$

Adding with (4.4), the total expected cost is at most $C + S \leq 2(1 + \alpha)C^* + (2 + \frac{2e^\alpha}{e^\alpha - 1}) \cdot S^*$. Since the optimal cost is $Z^* = C^* + S^*$, we choose $\alpha \approx 1.35$ to minimize the approximation ratio, and get the following result.

Theorem 4.3 *The algorithm SimpleCFL is a $\beta = 4.7$ -approximation algorithm for CFL.*

If we were to use an 1.55-approximation for Steiner tree [14] to buy edges in Step (S2), we would be getting an improved approximation ratio of 4.2 for CFL, while the analysis of [5] gets a 3.55 approximation. However, we need to use the 2-approximation algorithm of the MST, as there is a cross-monotone cost-sharing function ξ_{MST} for the MST problem.

4.2 A More Refined Assignment

We now present an improved 4.6-approximation. The idea behind the improvement is a natural one: instead of always going clockwise to look for the closest cluster D_l with a facility in it, we can look both clockwise and counter-clockwise for the closest cluster with a facility, breaking ties randomly. As one might expect, we show that this decreases the cost (see (4.17)), since it is no longer the case that a cluster with a facility in its neighboring counter-clockwise cluster goes several hops in the clockwise direction to find a (farther) facility.

More formally, let us modify the experiment of Section 4.1 to assign the demands to facilities in F as follows. Choose just one global ‘‘tie-breaker’’ bit TB at random. When $D_l \cap F = \emptyset$ and we go outside D_l , instead of always going around the cycle in a clockwise order, consider the smallest $t > 0$ such that either D_{l+t} or D_{l-t} has a non-empty intersection with F ; let D_s be the corresponding set with $D_s \cap F \neq \emptyset$. If both D_{l+t} and D_{l-t}

intersect F , then we break the tie randomly using TB: if $\text{TB} = 0$, then we choose D_{l-t} (i.e., set D_s to be D_{l-t}); else if $\text{TB} = 1$, we set D_s to be D_{l+t} . The same bit TB is used by all groups D_l . Note that $r \in D_0$, and so $t \leq \lceil k/2 \rceil$. We now send all the demands in D_l to the facility in D_s closest to f_s .

Let the random variables X_i and A_i be as in Section 4.1. We will not go over the analysis of this experiment in the same level of detail, but let us point out some of the salient ideas: for some facility in D_i to be used by $\geq \kappa$ groups of larger index (say), we must have: (i) $X_i = 0$, and (ii) either all 2κ groups $D_{i+1}, \dots, D_{i+2\kappa}$ do not intersect F , or only the last one does and we choose D_i with probability half. Like in (4.10), the expected total cost will again be a non-negative linear combination of the values $\{C'_i\}$ and $\{c(P'_i)\}$. Using the idea outlined in this paragraph, as well as expressions analogous to (4.8) and (4.9) in the previous analysis, we can verify that the coefficient of C'_i (in this linear combination) will be at most

$$1 + (1 - \mathbf{q}) + 2(1 - \mathbf{q}) \sum_{\kappa \geq 1} (\mathbf{q}^{2\kappa} + \frac{1}{2} \mathbf{q}^{2\kappa-1} (1 - \mathbf{q})) \leq 1 + (1 - \mathbf{q}) + \mathbf{q} = 2. \quad (4.13)$$

Analyzing the coefficient of $c(P'_i)$ needs more work. Assume without loss of generality that $\lfloor k/2 \rfloor \leq i \leq k$; the complementary case is similar. Given a set T of indices, let $U(T)$ denote the event that $X_j = 1$ for all $j \in T$; informally, the “ U ” denotes “unmarked”. We consider when the demands in group D_l will use the path P'_i ; we will do so by considering the cases $l \leq i$ and $l > i$.

Case I: $l \leq i$. Let \mathcal{A}_l be the event that the demands in group D_l use the path P'_i . For \mathcal{A}_l to hold, the following event \mathcal{B}_l should also happen: $\mathcal{B}_l \equiv U(\{j : |j - l| \leq i - l\})$. So,

$$\Pr[\mathcal{A}_l] = \Pr[\mathcal{B}_l] \times \Pr[\mathcal{A}_l \mid \mathcal{B}_l] = \mathbf{q}^{2(i-l)+1} \times \Pr[\mathcal{A}_l \mid \mathcal{B}_l].$$

Now, if $k + 1 - l \geq l$, it is easy to see by symmetry (since TB is equally likely to be 0 or 1) that $\Pr[\mathcal{A}_l \mid \mathcal{B}_l] \leq 1/2$; thus we have

$$\text{if } k + 1 - l \geq l, \text{ then } \Pr[\mathcal{A}_l] \leq \mathbf{q}^{2(i-l)+1}/2. \quad (4.14)$$

Next, suppose $k + 1 < 2l$. Define an event \mathcal{C}_l : $\mathcal{C}_l \equiv U(\{j : i + 1 - l \leq |j - l| \leq k - l\})$. We have $\Pr[\mathcal{A}_l \mid (\mathcal{B}_l \wedge \overline{\mathcal{C}}_l)] = 1/2$, by symmetry. Also, a moment's thought reveals that

$$\Pr[\mathcal{A}_l \mid (\mathcal{B}_l \wedge \mathcal{C}_l)] = 1 - (1 - \mathbf{q})/2 = (1 + \mathbf{q})/2.$$

Thus we have

$$\begin{aligned} \Pr[\mathcal{A}_l] &= \mathbf{q}^{2(i-l)+1} \cdot (\Pr[\overline{\mathcal{C}}_l \mid \mathcal{B}_l] \cdot (1/2) + \Pr[\mathcal{C}_l \mid \mathcal{B}_l] \cdot (1 + \mathbf{q})/2) \\ &= \mathbf{q}^{2(i-l)+1} \cdot ((1 - \mathbf{q}^{2(k-i)}) \cdot (1/2) + \mathbf{q}^{2(k-i)} \cdot (1 + \mathbf{q})/2) \\ &= \mathbf{q}^{2(i-l)+1} \cdot (1/2) \cdot (1 + \mathbf{q}^{2(k-i)+1}). \end{aligned}$$

Comparing with (4.14), we see that if $l \leq i$, then $\Pr[\mathcal{A}_l] \leq \mathbf{q}^{2(i-l)+1} \cdot (1/2) \cdot (1 + \mathbf{q}^{2(k-i)+1})$. Summing over all $l \leq i$, we get that the total contribution of all D_l , $l \leq i$, to the coefficient of $c(P'_i)$ in the expected connection cost, is at most

$$\frac{\mathbf{q}}{2(1 - \mathbf{q}^2)} \cdot (1 + \mathbf{q}^{2(k-i)+1}). \quad (4.15)$$

We now move to the next case.

Case II: $i < l \leq k$. Let \mathcal{A}_l be as in Case I. Now define $\mathcal{B}'_l \equiv U(\{j : |j - l| \leq l - (i + 1)\})$. Once again, \mathcal{B}'_l is a necessary condition for \mathcal{A}_l . Also note that $\Pr[\mathcal{B}'_l] = 0$ if $k + 1 - l \leq l - (i + 1)$; thus, we can restrict attention

to those $l > i$ such that $l \leq l_0 \doteq \lfloor (k+i+1)/2 \rfloor$. Now, since $l > i$, it is easy to see that $\Pr [\mathcal{A}_l \mid \mathcal{B}'_i] \leq 1/2$; also, $\Pr [\mathcal{B}'_i] = \mathbf{q}^{2(l-i)+1}$. Summing over all $l \in [i+1, l_0]$, we get a total contribution of

$$(1/2) \cdot \sum_{j=1}^{l_0-i} \mathbf{q}^{2j-1} \leq \frac{\mathbf{q}}{2(1-\mathbf{q}^2)} \cdot (1 - \mathbf{q}^{k-i+1}). \quad (4.16)$$

Adding with (4.16), the coefficient of $c(P'_i)$ in the expected connection cost is seen to be at most

$$\frac{\mathbf{q}}{2(1-\mathbf{q}^2)} \cdot (2 + \mathbf{q}^{2(k-i)+1} - \mathbf{q}^{k-i+1}) \leq \frac{\mathbf{q}}{1-\mathbf{q}^2} \leq \frac{e^{-\alpha}}{1-e^{-2\alpha}} = \frac{e^\alpha}{e^{2\alpha}-1}.$$

So, the total expected connection cost is at most $2C' + \frac{e^\alpha}{e^{2\alpha}-1} \cdot S'$. Using Lemma 4.2 to replace C' by $C^* + S^*$ and S' by $2S^*$, the expected connection cost is at most

$$2C^* + \frac{2(e^{2\alpha} + e^\alpha - 1)}{e^{2\alpha} - 1} \cdot S^*.$$

Adding with (4.4), the total expected cost is at most

$$2(1+\alpha)C^* + \frac{2(2e^{2\alpha} + e^\alpha - 2)}{e^{2\alpha} - 1} \cdot S^*. \quad (4.17)$$

Since the optimal cost is $Z^* = C^* + S^*$, we choose $\alpha = 1.296$ to minimize the approximation ratio to get the following result.

Theorem 4.4 *The algorithm SimpleCFL is a $\beta = 4.6$ -approximation algorithm for CFL.*

5 Analysis of the Limited-Independence Marking

We start with some useful derandomization tools in Section 5.1, and then analyze the limited-independence marking in Section 5.2.

5.1 Derandomization tools

We will use the following theorem, which appears as Lemma 2.3 in [2]:

Theorem 5.1 ([2]) *Let $t \geq 4$ be an even integer. Suppose X_1, \dots, X_n are t -wise independent random variables taking values in $[0, 1]$. Let $X = X_1 + \dots + X_n$ and $\mu = \mathbb{E}[X]$, and let $\lambda > 0$. Then*

$$\Pr[|X - \mu| \geq \lambda] \leq C_t \cdot \left(\frac{t\mu + t^2}{\lambda^2} \right)^{t/2},$$

where $C_t = 2\sqrt{\pi t} \cdot e^{1/(6t)} \cdot (5/(2e))^{t/2} \leq 8$.

See [15] for tail-bounds similar to the above.

We next present some notation.

- For each group D_j , let $Z_{j,1}, Z_{j,2}, \dots, Z_{j,M}$ be the respective indicator random variables for the demands in D_j getting marked by our algorithm, when these demands are considered in *nondecreasing order of distance from f_j* .
- Let A be any set of ordered pairs $\{(j, k)\}$. Then, $N(A)$ is the indicator random variable for the event “for all $(j, k) \in A$, $Z_{j,k} = 0$ ”. Also, $T(A)$ denotes $\sum_{(j,k) \in A} Z_{j,k}$. (“ N ” stands for “none”, and “ T ” for “total”.)

We now give two types of upper-bounds on $E[N(A)]$:

Lemma 5.2 *Let t_1 and t_2 be any even positive integers such that $t_1, t_2 \leq t$ (recall that the marking is done in a t -wise independent manner). The following hold for any set A of ordered pairs $\{(j, k)\}$.*

- (i) *Let $IE(s, A)$ be the random variable denoting the inclusion-exclusion expansion of $N(A)$ truncated at the s th level; i.e.,*

$$IE(s, A) = \sum_{r=0}^s (-1)^r \sum_{A' \subseteq A: |A'|=r} \prod_{(j,k) \in A'} Z_{j,k}.$$

Then, the inequality $N(A) \leq IE(t_1, A)$ holds always; also,

$$E[IE(t_1, A)] \leq (1 - \alpha/M)^{|A|} + \left(\frac{e\alpha|A|}{Mt_1} \right)^{t_1}. \quad (5.18)$$

- (ii) *Let $NCM(t_2, A)$ denote the “normalized central moment” $\frac{(T(A) - \alpha|A|/M)^{t_2}}{(\alpha|A|/M)^{t_2}}$. Then, the inequality $N(A) \leq NCM(t_2, A)$ holds always; also,*

$$\text{if } 4 \leq t_2 \leq \alpha|A|/M, \text{ then } E[NCM(t_2, A)] \leq 8 \cdot (2t_2)^{t_2/2} \cdot (\alpha|A|/M)^{-t_2/2}. \quad (5.19)$$

Proof: The upper bounds that are claimed to hold always on $N(A)$ in (i) and (ii), easily follow from the fact that t_1 and t_2 are even. Bound (5.18) follows from the statement and proof of Theorem 2 in [3]. Bound (5.19) follows from Theorem 5.1, using the fact that the “ $t\mu + t^2$ ” term in Theorem 5.1 is at most $2t\mu$ if $t \leq \mu$. ■

5.2 The analysis

Let ϵ be an arbitrary positive constant lying in $(0, 1)$. We now prove that if the demands are marked in t -wise independent fashion where $t = a \log(1/\epsilon)$ for a suitably large constant a , then the expected approximation ratio is at most $(1 + \epsilon)$ times what it is, in our analysis of independent marking in Section 4.2. The random bit (tie-breaker) TB introduced in Section 4.2 is still chosen independently of the marking; this only multiplies the size of our small sample space by a factor of 2.

By Lemma 4.1(a), we get that for the Steiner cost, we can continue to use the bound (4.4) even under t -wise independent marking; it is the connection cost that needs work. We now show that the expected total connection cost under the algorithm of Section 4.1 changes very little under limited-independence marking; we then briefly sketch the small modifications required for the algorithm of Section 4.2.

Recall the algorithm of Section 4.1. The total connection cost is a random variable that is the sum of three quantities: (i) the deterministic value $\sum_l C'_l$, which represents the cost of all demands in each D_l first getting routed to f_l ; (ii) the value that corresponds to unsuccessfully traveling through some D_l , and (iii) the total cost

paid in traveling from f_i to the closest marked demand in D_i , once D_i is identified as the closest cluster. We will only show that the expected total cost of (iii) gets multiplied by at most $(1 + \epsilon)$ due to our $t = a \log(1/\epsilon)$ -wise independent marking; the argument for the term (ii) is analogous, and, in fact, simpler. Specifically, we will show the following. Let i be the index of some arbitrary but fixed D_i . We show that the expected value of the random variable

$$\phi \doteq \sum_{j=1}^i M X_j X_{j+1} \cdots X_{i-1} \cdot (1 - X_i) A_i$$

gets multiplied by at most $(1 + \epsilon)$. (Since i is fixed, we have not subscripted ϕ as ϕ_i ; this remark also holds for some other definitions below.)

We now show how to use Lemma 5.2 to upper-bound $\mathbb{E}[\phi]$. Let a_1, a_2, \dots, a_M be the distances of the demands in D_i from f_i , written in *nondecreasing* order. Then, expanding the “ $(1 - X_i)A_i$ ” part of ϕ , we see that

$$\phi = \sum_{j=1}^i M X_j X_{j+1} \cdots X_{i-1} \cdot \left[\sum_{u=1}^M a_u Z_{i,u} \cdot \prod_{\ell=1}^{u-1} (1 - Z_{i,\ell}) \right].$$

Fix u arbitrarily, and let $z_u \doteq Z_{i,u} \cdot \prod_{\ell=1}^{u-1} (1 - Z_{i,\ell}) \cdot \sum_{j=1}^i X_j X_{j+1} \cdots X_{i-1}$. We aim to show that $\mathbb{E}[z_u]$ is multiplied by at most $(1 + \epsilon)$ in our t -wise independent marking, as compared to the fully-independent marking. For $j = 0, 1, \dots, i-1$, define $A_j = \{(r, s) : (i-j) \leq r \leq (i-1), 1 \leq s \leq M\}$. Thus we get

$$z_u = Z_{i,u} \cdot N(\{(i, \ell) : 1 \leq \ell \leq u-1\}) \cdot \sum_{j=0}^{i-1} N(A_j).$$

Now let a_1 be a sufficiently large constant. Define a_2 to be the smallest even integer that is at least $2e \log(1/\epsilon)$, t_1 to be the smallest even integer that is at least $2ea_1 \log(1/\epsilon)$, and t_2 to be the largest even integer that is at most $a_1 \log(1/\epsilon)/4$. There are two cases: $a_1 \log(1/\epsilon) \leq i-1$, or $a_1 \log(1/\epsilon) > i-1$; we start with the harder first case.

Case I: $a_1 \log(1/\epsilon) \leq i-1$. Break up the expression for z_u into two sums, one for “small” j and the other for “larger” j :

$$Z_{i,u} \cdot N(\{(i, \ell) : 1 \leq \ell \leq u-1\}) \cdot \sum_{j \leq a_1 \log(1/\epsilon)} N(A_j) + Z_{i,u} \cdot N(\{(i, \ell) : 1 \leq \ell \leq u-1\}) \cdot \sum_{j > a_1 \log(1/\epsilon)} N(A_j).$$

Thus, by Lemma 5.2, z_u is always bounded by the sum of the following two random variables:

$$Z_{i,u} \cdot IE(a_2, \{(i, \ell) : 1 \leq \ell \leq u-1\}) \cdot \sum_{j \leq a_1 \log(1/\epsilon)} IE(t_1, A_j), \text{ and} \quad (5.20)$$

$$Z_{i,u} \cdot IE(a_2, \{(i, \ell) : 1 \leq \ell \leq u-1\}) \cdot \sum_{j > a_1 \log(1/\epsilon)} NCM(t_2, A_j). \quad (5.21)$$

If we expand the expectations of these two random variables using the linearity of expectation, we get terms each of which is a product of some of the random variables $Z_{i,\cdot}$; importantly, the number of factors in each such term is at most $1 + a_2 + t_1$ and $1 + a_2 + t_2$ respectively. Thus, if we choose $t = 1 + a_2 + t_1$ (recall that $t_1 \geq t_2$), then the expectations of these two random variables become

$$\mathbb{E}[Z_{i,u}] \cdot \mathbb{E}[IE(a_2, \{(i, \ell) : 1 \leq \ell \leq u-1\})] \cdot \sum_{0 \leq j \leq a_1 \log(1/\epsilon)} \mathbb{E}[IE(t_1, A_j)] \text{ and} \quad (5.22)$$

$$\mathbb{E}[Z_{i,u}] \cdot \mathbb{E}[IE(a_2, \{(i, \ell) : 1 \leq \ell \leq u-1\})] \cdot \sum_{j > a_1 \log(1/\epsilon)} \mathbb{E}[NCM(t_2, A_j)] \quad (5.23)$$

respectively. We next use (5.18) and (5.19) to bound these values; we will see that choosing the constant a_1 large enough results in $\mathbb{E}[z_u]$ being at most $(1 + \epsilon)$ times what it is with independent marking.

The expression (5.22) is at most

$$\frac{\alpha}{M} \cdot \left((1 - \alpha/M)^{u-1} + \left(\frac{e\alpha(u-1)}{Ma_2} \right)^{a_2} \right) \cdot \sum_{0 \leq j \leq a_1 \log(1/\epsilon)} \left((1 - \alpha/M)^{Mj} + (e\alpha j/t_1)^{t_1} \right);$$

i.e., at most

$$\frac{\alpha}{M} \cdot \left((1 - \alpha/M)^{u-1} + (e\alpha/a_2)^{a_2} \right) \cdot \sum_{0 \leq j \leq a_1 \log(1/\epsilon)} \left((1 - \alpha/M)^{Mj} + (e\alpha a_1 \log(1/\epsilon)/t_1)^{t_1} \right).$$

Similarly, the expression (5.23) is at most

$$\frac{\alpha}{M} \cdot \left((1 - \alpha/M)^{u-1} + (e\alpha/a_2)^{a_2} \right) \cdot \sum_{j > a_1 \log(1/\epsilon)} \left(8 \cdot (2t_2/(\alpha j))^{t_2/2} \right).$$

Thus, if the demands are marked in t -wise independent fashion, then $\mathbb{E}[z_u]$ is at most $(\alpha/M) \cdot \left((1 - \alpha/M)^{u-1} + (e\alpha/a_2)^{a_2} \right)$ times

$$\sum_{0 \leq j \leq a_1 \log(1/\epsilon)} \left((1 - \alpha/M)^{Mj} + (e\alpha a_1 \log(1/\epsilon)/t_1)^{t_1} \right) + \sum_{j > a_1 \log(1/\epsilon)} \left(8 \cdot (2t_2/(\alpha j))^{t_2/2} \right).$$

On the other hand, under fully-independent marking,

$$\mathbb{E}[z_u] = \frac{\alpha}{M} \cdot (1 - \alpha/M)^{u-1} \cdot \sum_{j=0}^{i-1} (1 - \alpha/M)^{Mj}. \quad (5.24)$$

Recalling the definitions of a_1 , a_2 , t_1 and t_2 , it is easy to verify that if a_1 is chosen as a sufficiently large constant, then the former value is at most $(1 + \epsilon)$ times the latter. This completes the analysis of Case I.

Case II: $a_1 \log(1/\epsilon) > i - 1$. We follow the above analysis; we need not consider the case of larger j . Now, under limited-independence marking, the bound on $\mathbb{E}[z_u]$ is simpler: it is at most

$$(\alpha/M) \cdot \left((1 - \alpha/M)^{u-1} + (e\alpha/a_2)^{a_2} \right) \cdot \sum_{0 \leq j \leq i-1} \left((1 - \alpha/M)^{Mj} + (e\alpha a_1 \log(1/\epsilon)/t_1)^{t_1} \right).$$

Once again, this is at most $(1 + \epsilon)$ times the r.h.s. of (5.24).

Modifications required for the algorithm of Section 4.2. We now sketch the few modifications required. Conditional on $TB = 1$, the demands in D_l will get routed to:

- D_l , iff $X_l = 0$;
- D_{l+t} for some $t > 0$, iff $X_l(1 - X_{l+t}) \prod_{j=1}^{t-1} (X_{l+j} X_{l-j})$ holds; and to

- D_{l-t} for some $t > 0$, iff $X_l X_{l+t} (1 - X_{l-t}) \prod_{j=1}^{t-1} (X_{l+j} X_{l-j})$ holds.

Similar remarks hold conditional on $TB = 0$. The expected connection cost is $1/2$ times the cost conditional on $TB = 1$, plus $1/2$ times the cost conditional on $TB = 0$. Consider the case where $TB = 1$. The total connection cost can then be verified to still be a non-negative linear combination of terms of the form: “a product of some X_j ’s, multiplied by at most one term of the form $(1 - X_l)A_l$ ”. Furthermore, it remains true of the total connection cost that the expected values of: (a) the coefficient of any $c(P_i')$, and (b) the total term multiplying any term of the form $(1 - X_l)A_l$, are both $\Theta(M)$. Using these observations, the above type of analysis can be seen to hold for the algorithm of Section 4.2 also.

6 Further Extensions

6.1 Handling Facility Costs

For the model of connected facility location where there are facility opening costs as well, we can run a slight variant of the algorithm above:

- F1. *Mark* each demand $j \in D$ independently with probability $1/M$, and let D' be the set of marked demands.
- F2. Compute a ρ_{FL} -approximate metric facility location solution with the original distances and facility opening costs, but where the demand value for any $j \in D'$ is M , and is 0 for $j \notin D'$. Let F be the set of open facilities returned by this algorithm.
- F3. Construct a ρ_{ST} -approximate Steiner tree T on $F \cup \{r\}$.
- F4. Assign each demand $j \in D$ to its closest open facility $i(j)$ in F .

The cost analysis is very similar to that of Theorem 2, and we only sketch the important ideas here. Let O^* , C^* and S^* denote the facility opening costs, connection costs and Steiner costs of the optimal solution. Note that opening the same set of facilities as OPT is also a feasible solution to the facility location instance in Step (F2) with expected cost $(O^* + C^*)$, and hence we pay

$$A_1 \doteq \rho_{FL}(O^* + C^*) \tag{6.25}$$

in expectation, where ρ_{FL} is the approximation guarantee of the algorithm used in Step (F2). Now a proof along the lines of Lemma 4.1 implies that there is a Steiner tree connecting these facilities with expected cost $(S^* + C^* + A_1)$, and hence the tree we construct in Step F3 costs at most ρ_{ST} times that, which is

$$A_2 \doteq \rho_{ST}(S^* + C^* + \rho_{FL}(C^* + O^*)). \tag{6.26}$$

Finally, we have to bound the connection costs of Step (F4): one can use an analysis very similar to Section 4.2, but in addition to the expected cost of $2C^* + \frac{2e}{e-1}S^*$ paid there, one also has to pay an additional cost to go from the marked demands in D' to the facilities in F : this costs A_1 in expectation. Hence the expected assignment cost in Step (F4) is

$$A_3 \doteq 2C^* + \frac{2e}{e-1}S^* + (C^* + O^*)\rho_{FL}. \tag{6.27}$$

Adding up the three expressions above, we get a bound on the expected cost of at most

$$\rho_{FL}(2 + \rho_{ST})O^* + (1 + \rho_{FL})(2 + \rho_{ST})C^* + (\rho_{ST} + 2e/(e - 1))S^*. \tag{6.28}$$

Now using the best known approximation factors of $\rho_{FL} = 1.52$ [12] and $\rho_{ST} = 1.55$ [14], we get a 8.94-approximate algorithm for the case of non-zero cost for opening facilities. (A better performance can be obtained by choosing α carefully.)

To get cross-monotonic cost shares, we can instead use the $\rho_{FL} = 3$ approximate Facility Location algorithm of Pál and Tardos [13] and its accompanying cross-monotone cost sharing scheme ξ_{FL} , the MST heuristic with $\rho_{ST} = 2$ for Steiner tree (and use its cross-monotone cost-sharing scheme ξ_{MST} as well). This gives us an approximation factor of $\beta = 16$; now setting the cost-share of each demand j to be

$$\xi(D, j) = \frac{1}{\beta} \mathbb{E}[M \xi_{MST}(D' \cup r, j) + M \xi_{FL}(D', j) + \ell(j, D' \cup r)], \quad (6.29)$$

can give us a β -approximate cross-monotone cost sharing scheme for CFL with facility costs as well.

6.2 Handling Non-Uniform Demands

The discussion in the previous sections assumed that each of the demands $j \in D$ had a weight $d_j = 1$; i.e., it wanted to send just one unit of demand to the root. Moreover, we assumed that there was only one player at each vertex in D . Let us briefly indicate how these assumptions can be discharged; in this section, we do not attempt to optimize the constants, sacrificing the performance guarantees in favor of sketching the main ideas.

Let us note that handling the assumption of a single player at every vertex is simple: if there were n_v players at vertex v , we create n_v new vertices, attach each of them to v using zero-cost edges, and place a unique player from v on each of the new vertices.

To handle arbitrary weights d_j for demand j , we first take care of the following two special cases separately, which ensure that the values of d_j all lie within a multiplicative factor of n^2 of each other.

Heavy Vertices If the weight d_j of vertex j is at least M , we can *deterministically mark* the vertex j in Step S1.

It can be shown that OPT would have connected these vertices up in its Steiner tree as well, and hence this causes no loss in quality.

Light Vertices If the weight d_j of node j is at most M/n^2 , then we do not mark j at all in Step S1: note that the total weight $\sum_{j \in V} d_j$ of such nodes is at most $n \cdot M/n^2 = M/n$. The expected cost of the Steiner tree can only go down due to this change, and a slight variant on the proof of Theorem 4.4 will show that the expected assignment cost only increases by a factor of $(1 + 1/n)$ —essentially, throwing back the demand of M/n in any subtree can only cause an increase of a factor of $(1 + 1/n)$ in the expected assignment cost.

Having performed these operations, we can now rescale and assume that the weights d_j lie in $[n, n^3]$, and that the parameter M is greater than n^3 . We round the weights d_j down to the closest integer $\lfloor d_j \rfloor$: this decreases each weight by at most a $(1 + 1/n)$ factor, and hence alters only the lower order terms of our performance guarantee. And finally, we replace the player at j by $\lfloor d_j \rfloor$ players: a long but straight-forward reworking of the proofs in Section 4 shows that the algorithm is a constant-factor approximation even with general weights, and gives us the desired cost-shares.

Acknowledgment. We thank the referees for their helpful comments.

References

- [1] Noga Alon, László Babai, and Alon Itai. A fast and simple randomized parallel algorithm for the maximal independent set problem. *Journal of Algorithms*, 7:567–583, 1986.
- [2] Mihir Bellare and John Rompel. Randomness-efficient oblivious sampling. In *Proc. IEEE Symposium on Foundations of Computer Science*, pages 276–287, 1994.
- [3] Guy Even, Oded Goldreich, Michael Luby, Noam Nisan, and Boban Veličković. Approximations of general independent distributions. In *Proceedings of the 24th Annual ACM Symposium on Theory of Computing*, pages 10–16, 1992.
- [4] Anupam Gupta, Amit Kumar, Jon Kleinberg, Rajeev Rastogi, and Bülent Yener. Provisioning a Virtual Private Network: A network design problem for multicommodity flow. In *Proceedings of the 33rd Annual ACM Symposium on Theory of Computing*, pages 389–398, 2001.
- [5] Anupam Gupta, Amit Kumar, and Tim Roughgarden. Simpler and better approximation algorithms for network design. In *Proceedings of the 35th Annual ACM Symposium on Theory of Computing*, pages 365–372, 2003.
- [6] Kathryn J. Kent and Darko Skorin-Kapov. Population monotonic cost allocations on MSTs. In *Proceedings of the 6th International Conference on Operational Research (Rovinj, 1996)*, pages 43–48. Croatian Oper. Res. Soc., Zagreb, 1996.
- [7] Michael Luby. A simple parallel algorithm for the maximal independent set problem. *SIAM J. Comput.*, 15(4):1036–1053, 1986.
- [8] Hervé Moulin and Scott Shenker. Strategyproof sharing of submodular costs: budget balance versus efficiency. *Economic Theory* 18, pp. 511–533, 2001.
- [9] Kamal Jain and Vijay Vazirani. Applications of approximation algorithms to cooperative games. In *Proceedings of the 33rd Annual ACM Symposium on the Theory of Computing (STOC)*, pages 364–372, 2001.
- [10] David R. Karger and Maria Minkoff. Building Steiner trees with incomplete global knowledge. In *Proceedings of the 41th Annual IEEE Symposium on Foundations of Computer Science*, pages 613–623, 2000.
- [11] Stefano Leonardi and Guido Schäfer. Cross-monotonic cost-sharing methods for connected facility location games. *Theoretical Computer Science* **326**(1-3), 431–442, 2004.
- [12] Mohammad Mahdian, Yinyu Ye, and Jiawei Zhang. Improved approximation algorithms for metric facility location problems. In *Approximation algorithms for combinatorial optimization*, volume 2462 of *Lecture Notes in Comput. Sci.*, pages 229–242. Springer, Berlin, 2002.
- [13] Martin Pál and Éva Tardos. Group Strategyproof Mechanisms via Primal-Dual Algorithms. In *Proceedings of the 44th Annual IEEE Symposium on Foundations of Computer Science*, 2003, pp. 584–593.
- [14] Gabriel Robins and Alexander Zelikovsky. Improved Steiner tree approximation in graphs. In *Proceedings of the 11th Annual ACM-SIAM Symposium on Discrete Algorithms*, 2000, pp. 770–779.
- [15] Jeanette P. Schmidt, Alan Siegel, and Aravind Srinivasan. Chernoff-Hoeffding bounds for applications with limited independence, *SIAM J. Discrete Math.*, 8 (1995), pp. 223–250.

- [16] H. P. Young. Cost allocation. In R. J. Aumann and S. Hart, editors, *Handbook of Game Theory*, volume 2, chapter 34, pages 1193–1235. North-Holland, 1994.