

Machine Learning


10-701/15-781, Spring 2008

Computational Learning Theory

Eric Xing

Lecture 11, February 20, 2008

Reading: Chap. 7 T.M book



Complexity of Learning

- The complexity of learning is measured mainly along two axis: **Information** and **computation**.

The **Information complexity** is concerned with the generalization performance of learning;

- How many training examples are needed?*
- How fast do learner's estimate converge to the true population parameters? Etc.*

The **Computational complexity** concerns the computation resources applied to the training data to extract from it learner's predictions.

It seems that when an algorithm improves with respect to one of these measures it deteriorates with respect to the other.



What General Laws constrain Inductive Learning?



- Sample Complexity
 - How many training examples are sufficient to learn target concept?
- Computational Complexity
 - Resources required to learn target concept?
- Want theory to relate:
 - Training examples
 - Quantity m
 - Quality
 - How presented
 - Complexity of hypothesis/concept space H
 - Accuracy of approx to target concept ϵ
 - Probability of successful learning δ

These results only useful wrt $O(\dots)$!

Prototypical concept learning task



Binary classification

- Everything we'll say here generalizes to other, including regression and multi-class classification, problems.
- Given:
 - Instances X : Possible days, each described by the attributes *Sky, AirTemp, Humidity, Wind, Water, Forecast*
 - Target function c : *EnjoySport* : $X \rightarrow \{0, 1\}$
 - Hypotheses space H : Conjunctions of literals. E.g.
 $(?, \text{Cold}, \text{High}, ?, ?, \text{EnjoySport})$.
 - Training examples S : iid positive and negative examples of the target function
 $(x_1, c(x_1)), \dots (x_m, c(x_m))$
- Determine:
 - A hypothesis h in H such that $h(x)$ is "good" w.r.t $c(x)$ for all x in S ?
 - A hypothesis h in H such that $h(x)$ is "good" w.r.t $c(x)$ for all x in the true dist \mathcal{D} ?

Two Basic Competing Models



PAC framework

Sample labels are consistent with some h in H

Learner's hypothesis required to meet *absolute* upper bound on its error

Agnostic framework

No prior restriction on the sample labels

The required upper bound on the hypothesis error is only relative (to the best hypothesis in the class)


Sample Complexity




- How many training examples are sufficient to learn the target concept?
- Training scenarios:
 - 1 If learner proposes instances, as queries to teacher
 - Learner proposes instance x , teacher provides $c(x)$
 - 2 If teacher (who knows c) provides training examples
 - teacher provides sequence of examples of form $(x, c(x))$
 - 3 If some random process (e.g., nature) proposes instances
 - instance x generated randomly, teacher provides $c(x)$

Protocol

- Given:
 - set of examples X
 - fixed (unknown) distribution D over X
 - set of hypotheses H
 - set of possible target concepts C
- Learner observes sample $S = \{ \langle x_i, c(x_i) \rangle \}$
 - instances x_i drawn from distr. D
 - labeled by target concept $c \in C$
(Learner does NOT know $c(\cdot), D$)
- Learner outputs $h \in H$ estimating c
 - h is evaluated by performance on subsequent instances drawn from D
- For now:
 - $C = H$ (so $c \in H$)
 - Noise-free data



True error of a hypothesis



- Definition: The **true error** (denoted $\epsilon_D(h)$) of hypothesis h with respect to target concept c and distribution \mathcal{D} is the probability that h will misclassify an instance drawn at random according to \mathcal{D} .

$$\epsilon_D(h) \equiv \Pr_{x \in \mathcal{D}}[c(x) \neq h(x)]$$

Two notions of error



- *Training error* of hypothesis h with respect to target concept c
 - How often $h(x) \neq c(x)$ over training instance from \mathcal{S}

$$\hat{\epsilon}_{\mathcal{S}}(h) \equiv \Pr_{x \in \mathcal{S}}[c(x) \neq h(x)] \equiv \frac{\sum_{x \in \mathcal{S}} \delta(c(x) \neq h(x))}{|\mathcal{S}|}$$

- *True error* of hypothesis h with respect to c
 - How often $h(x) \neq c(x)$ over future random instances drawn iid from \mathcal{D}

$$\epsilon_{\mathcal{D}}(h) \equiv \Pr_{x \in \mathcal{D}}[c(x) \neq h(x)]$$

Can we bound
 $\hat{\epsilon}_{\mathcal{S}}(h)$
in terms of
 $\hat{\epsilon}_{\mathcal{D}}(h)$
??

The Union Bound



- Lemma. (The union bound). Let A_1, A_2, \dots, A_k be k different events (that may not be independent). Then

$$P(A_1 \cup A_2 \cup \dots \cup A_k) \leq P(A_1) + P(A_2) + \dots + P(A_k)$$

- In probability theory, the union bound is usually stated as an axiom (and thus we won't try to prove it), but it also makes intuitive sense: The probability of any one of k events happening is at most the sum of the probabilities of the k different events.

Hoeffding inequality



- Lemma. (Hoeffding inequality) Let Z_1, \dots, Z_m be m independent and identically distributed (iid) random variables drawn from a Bernoulli(ϕ) distribution, i.e., $P(Z_i = 1) = \phi$, and $P(Z_i = 0) = 1 - \phi$.

Let $\hat{\phi} = (1/m) \sum_{i=1}^m Z_i$ be the mean of these random variables, and let any $\gamma > 0$ be fixed. Then

$$P(|\phi - \hat{\phi}| > \gamma) \leq 2 \exp(-2\gamma^2 m)$$

- This lemma (which in learning theory is also called the Chernoff bound) says that if we take $\hat{\phi}$ — the average of m Bernoulli(ϕ) random variables — to be our estimate of ϕ , then the probability of our being far from the true value is small, so long as m is large.

Version Space




- A hypothesis h is consistent with a set of training examples \mathcal{S} of target concept c if and only if $h(x) = c(x)$ for each training example $\langle x_i, c(x_i) \rangle$ in \mathcal{S}

$$\text{Consistent}(h, \mathcal{S}) \models h(x) = c(x), \forall \langle x, c(x) \rangle \in \mathcal{S}$$

- The version space, $VS_{H, \mathcal{S}}$, with respect to hypothesis space H and training examples \mathcal{S} is the subset of hypotheses from H consistent with all training examples in \mathcal{S} .

$$VS_{H, \mathcal{S}} \equiv \{h \in H \mid \text{Consistent}(h, \mathcal{S})\}$$

Exhausting the version space



(r = training error, $error$ = true error)

- Definition: The version space $VS_{H,S}$ is said to be ϵ -exhausted with respect to c and S , if every hypothesis h in $VS_{H,S}$ has error less than ϵ with respect to c and S .

$$\forall h \in VS_{H,S}, \quad \hat{\epsilon}_S(h) < \epsilon$$

Probably Approximately Correct



Goal:

PAC-Learner produces hypothesis \hat{h} that

is approximately correct,

$$err_D(\hat{h}) \approx 0$$

with high probability

$$P(err_D(\hat{h}) \approx 0) \approx 1$$

- Double "hedging"

- approximately

- probably

Need both!

How many examples will ϵ -exhaust the VS



Theorem: [Haussler, 1988].

- If the hypothesis space H is finite, and S is a sequence of $m \geq 1$ independent random examples of some target concept c , then for any $0 \leq \epsilon \leq 1$, the probability that the version space with respect to H and D is not ϵ -exhausted (with respect to c) is less than

$$|H|e^{-\epsilon m}$$

- This bounds the probability that any consistent learner will output a hypothesis h with $\epsilon(h) \geq \epsilon$
- If we want to this probability to be below δ

$$|H|e^{-\epsilon m} \leq \delta$$

- then

$$m \geq \frac{1}{\epsilon} (\ln |H| + \ln(1/\delta))$$

Proof



What it means



- [Haussler, 1988]: probability that the version space is not ϵ -exhausted after m training examples is at most $|H|e^{-\epsilon m}$

$$Pr(\exists h \in H, \text{ s.t. } (error_{train}(h) = 0) \wedge (error_{true}(h) > \epsilon)) \leq |H|e^{-\epsilon m}$$

Suppose we want this probability to be at most δ

1. How many training examples suffice?

$$m \geq \frac{1}{\epsilon} (\ln |H| + \ln(1/\delta))$$

2. If $error_{train}(h) = 0$ then with probability at least $(1-\delta)$:

$$error_{true} \leq \frac{1}{m} (\ln |H| + \ln(1/\delta))$$

Learning Conjunctions of Boolean Literals



- How many examples are sufficient to assure with probability at least $(1 - \delta)$ that

every h in $VS_{H,S}$ satisfies $\epsilon_S(h) \leq \epsilon$

- Use our theorem:

$$m \geq \frac{1}{\epsilon} (\ln |H| + \ln(1/\delta))$$

- Suppose H contains conjunctions of constraints on up to n boolean attributes (i.e., n boolean literals).

Then $|H| = 3^n$, and

$$m \geq \frac{1}{\epsilon} (\ln 3^n + \ln(1/\delta))$$

or

$$m \geq \frac{1}{\epsilon} (n \ln 3 + \ln(1/\delta))$$

How about *EnjoySport*?



$$m \geq \frac{1}{\epsilon} (\ln |H| + \ln(1/\delta))$$

- If H is as given in *EnjoySport* then $|H| = 973$, and

$$m \geq \frac{1}{\epsilon} (\ln 973 + \ln(1/\delta))$$

- if want to assure that with probability 95%, VS contains only hypotheses with $\epsilon_S(h) \leq .1$, then it is sufficient to have m examples, where

$$m \geq \frac{1}{.1} (\ln 973 + \ln(1/.05))$$

$$m \geq 10(\ln 973 + \ln 20)$$

$$m \geq 10(6.88 + 3.00)$$

$$m \geq 98.8$$

PAC-Learning



- Learner L can draw labeled instance $\langle x, c(x) \rangle$ in unit time, $x \in X$ of length n drawn from distribution \mathcal{D} , labeled by target concept $c \in C$

Def'n: Learner L PAC-learns class C using hypothesis space H

if

1. for any target concept $c \in C$,
any distribution \mathcal{D} , any ϵ such that $0 < \epsilon < 1/2$, δ such that $0 < \delta < 1/2$,
 L returns $h \in H$ s.t.
w/ prob. $\geq 1 - \delta$, $\text{err}_{\mathcal{D}}(h) < \epsilon$
2. L 's run-time (and hence, sample complexity)
is $\text{poly}(|x|, \text{size}(c), 1/\epsilon, 1/\delta)$

- Sufficient:

1. Only $\text{poly}(\dots)$ training instances — $|H| = 2^{\text{poly}()}$
2. Only poly time / instance ...

Often $C = H$

Agnostic Learning



So far, assumed $c \in H$

Agnostic learning setting: don't assume $c \in H$

- What do we want then?
 - The hypothesis h that makes fewest errors on training data
- What is sample complexity in this case?

$$m \geq \frac{1}{2\varepsilon^2} (\ln|H| + \ln(1/\delta))$$

derived from Hoeffding bounds:

$$\Pr[\text{error}_D(h) > \text{error}_S(h) + \varepsilon] \leq e^{-2m\varepsilon^2}$$

Empirical Risk Minimization Paradigm



- Choose a **Hypothesis Class** H of subsets of X .
- For an input sample S , find some h in H that fits S "well".
- For a new point x , predict a label according to its membership in h .

$$\hat{h} = \arg \min_{h \in H} \hat{\varepsilon}_S(h)$$

- Example:
 - Consider linear classification, and let $h_\theta(x) = 1\{\theta^T x \geq 0\}$
Then $H = \{h_\theta : h_\theta(x) = 1\{\theta^T x \geq 0\}, \theta \in R^{n+1}\}$

$$\hat{\theta} = \arg \min_{\theta} \hat{\varepsilon}_S(h_\theta)$$

- We think of ERM as the most "basic" learning algorithm, and it will be this algorithm that we focus on in the remaining.
- In our study of learning theory, it will be useful to abstract away from the specific parameterization of hypotheses and from issues such as whether we're using a linear classifier or an ANN



The Case of Finite H

- $H = \{h_1, \dots, h_k\}$ consisting of k hypotheses.
- We would like to give guarantees on the generalization error of \hat{h} .
- First, we will show that $\hat{\epsilon}(h)$ is a reliable estimate of $\epsilon(h)$ for all h .
- Second, we will show that this implies an upper-bound on the generalization error of \hat{h} .



Misclassification Probability

- The outcome of a binary classifier can be viewed as a Bernoulli random variable $Z : Z = 1\{h_i(x) \neq c(x)\}$
- For each sample: $Z_j = 1\{h_i(x_j) \neq c(x_j)\}$

$$\hat{\epsilon}(h_i) = \frac{1}{m} \sum_{j=1}^m Z_j$$

- Hoeffding inequality

$$P(|\epsilon(h_i) - \hat{\epsilon}(h_i)| > \gamma) \leq 2\exp^{-2\gamma^2 m}$$

- This shows that, for our particular h_p , training error will be close to generalization error with high probability, assuming m is large.

Uniform Convergence



- But we don't just want to guarantee that $\hat{\epsilon}(h_i)$ will be close $\epsilon(h_i)$ (with high probability) for just only one particular h_i . We want to prove that this will be true for simultaneously for all $h_i \in H$
- For k hypothesis:

$$\begin{aligned}
 P(\exists h \in H, |\epsilon(h_i) - \hat{\epsilon}(h_i)| > \gamma) &= P(A_1 \cup \dots \cup A_k) \\
 &= \sum_{i=1}^k P(A_i) \\
 &= \sum_{i=1}^k 2 \exp(-2\gamma^2 m) \\
 &= 2k \exp(-2\gamma^2 m)
 \end{aligned}$$

- This means:

$$\begin{aligned}
 P(\neg \exists h \in H, |\epsilon(h_i) - \hat{\epsilon}(h_i)| > \gamma) &= P(\forall h \in H, |\epsilon(h_i) - \hat{\epsilon}(h_i)| \leq \gamma) \\
 &= 1 - 2k \exp(-2\gamma^2 m)
 \end{aligned}$$



- In the discussion above, what we did was, for particular values of m and γ , given a bound on the probability that:

for some $h_i \in H$

$$|\epsilon(h_i) - \hat{\epsilon}(h_i)| > \gamma$$

- There are three quantities of interest here: m and γ , and probability of error; we can bound either one in terms of the other two.

Sample Complexity



- How many training examples we need in order make a guarantee?

$$P(\exists h \in H, |\epsilon(h) - \hat{\epsilon}(h)| > \gamma) = 2k \exp(-2\gamma^2 m)$$

- We find that if $m \geq \frac{1}{2\gamma^2} \log \frac{2k}{\delta}$

then with probability at least $1-\delta$, we have that $|\epsilon(h_i) - \hat{\epsilon}(h_i)| \leq \gamma$
for all $h_i \in H$

- The key property of the bound above is that the number of training examples needed to make this guarantee is only logarithmic in k , the number of hypotheses in H . This will be important later.

Generalization Error Bound



- Similarly, we can also hold m and δ fixed and solve for γ in the previous equation, and show [again, convince yourself that this is right!] that with probability $1-\delta$, we have that for all $h_i \in H$

$$|\hat{\epsilon}(h) - \epsilon(h)| \leq \sqrt{\frac{1}{m} \log \frac{2k}{\delta}}$$

- Define $h^* = \arg \min_{h \in H} \epsilon(h)$ to be the best possible hypothesis in H .

$$\begin{aligned} \epsilon(\hat{h}) &\leq \hat{\epsilon}(\hat{h}) + \gamma \\ &\leq \hat{\epsilon}(\hat{h}^*) + \gamma \\ &\leq \epsilon(\hat{h}^*) + \gamma \end{aligned}$$

- If uniform convergence occurs, then the generalization error of $\hat{\epsilon}(h)$ is at most 2γ worse than the best possible hypothesis in H !

Summary



Theorem. Let $|\mathcal{H}| = k$, and let any m, δ be fixed. Then with probability at least $1 - \delta$, we have that

$$\varepsilon(\hat{h}) \leq \left(\min_{h \in \mathcal{H}} \varepsilon(h) \right) + 2\sqrt{\frac{1}{2m} \log \frac{2k}{\delta}}.$$

Corollary. Let $|\mathcal{H}| = k$, and let any δ, γ be fixed. Then for $\varepsilon(\hat{h}) \leq \min_{h \in \mathcal{H}} \varepsilon(h) + 2\gamma$ to hold with probability at least $1 - \delta$, it suffices that

$$\begin{aligned} m &\geq \frac{1}{2\gamma^2} \log \frac{2k}{\delta} \\ &= O\left(\frac{1}{\gamma^2} \log \frac{k}{\delta}\right), \end{aligned}$$