

How to (Re)Invent Tait’s Method*

Robert Harper

Spring Semester, 2022

1 Introduction

Two of the most important developments in type theory were the invention, by W. W. Tait, of *Tait’s Method* for function types, which was later extended by *J.Y.Girard* to *Girard’s Method* for type quantification, both of which were incorporated into a general theory of *logical relations* for a wide range of type theories. Tait’s method continues to be known by its original name, the *computability method*, which interprets types as predicates in the manner developed below.

The problem considered by Tait was to prove that β -reduction for the simply typed λ -calculus is *strongly normalizing*, which is usually defined to mean that there are *no* infinite β -reduction sequences starting with a well-typed term: $M = M_0 \rightarrow_{\beta} M_1 \rightarrow \dots$. A better definition, of more immediate utility, is the validity of *transfinite induction on reduction*, stated as follows: for any property \mathcal{P} of typed λ -terms, to show that \mathcal{P} holds for every such term, it is enough to show, for every well-typed typed term M , if all of its immediate β -reducts satisfy \mathcal{P} , then so does \mathcal{P} . More succinctly,

$$(\forall M:\tau(\forall N:\tau.M \rightarrow_{\beta} N \supset \mathcal{P}(N)) \supset \mathcal{P}(M)) \supset \forall M:\tau.\mathcal{P}(M).$$

The importance of strong normalization lies exactly in the utility afforded by this principle for proving other properties. For example, using transfinite induction on reduction it is possible to prove that weak confluence (every one-step split can be reconciled) implies confluence (transitivity of the “has a common reduct” relation.) This statement is false for untyped terms; it is precisely the strong normalization property that ensures its validity for well-typed terms.

The question considered here is a related, but technically much simpler, problem, the termination of a deterministic head reduction strategy for a simply typed λ -calculus. The simplification compared to Tait’s original proof is that head reduction is defined only for closed terms, avoiding the need to consider open terms in the development of Tait’s method. The type system considered here has unit, product, and function types, augmented with a two-element type of *observables*, corresponding to the familiar “accept/reject” formulation in the study of abstract machines.

*Copyright © Robert Harper. All Rights Reserved.

$$\begin{array}{c}
\text{VAR} \qquad \qquad \text{YES} \qquad \qquad \text{NO} \qquad \qquad \text{UNIT} \qquad \qquad \text{PAIR} \\
\frac{}{\Gamma, x:A \vdash x : A} \qquad \frac{}{\Gamma \vdash \text{yes} : \mathbf{2}} \qquad \frac{}{\Gamma \vdash \text{no} : \mathbf{2}} \qquad \frac{}{\Gamma \vdash \star : \mathbf{1}} \qquad \frac{\Gamma \vdash M_1 : A_1 \quad \Gamma \vdash M_2 : A_2}{\Gamma \vdash \langle M_1, M_2 \rangle : A_1 \times A_2} \\
\\
\text{LFT} \qquad \qquad \text{RHT} \qquad \qquad \text{LAM} \\
\frac{\Gamma \vdash M : A_1 \times A_2}{\Gamma \vdash M \cdot \mathbf{1} : A_1} \qquad \frac{\Gamma \vdash M : A_1 \times A_2}{\Gamma \vdash M \cdot \mathbf{2} : A_2} \qquad \frac{\Gamma, x:A_1 \vdash M_2 : A_2}{\Gamma \vdash \lambda_{A_1}(x . M_2) : A_1 \rightarrow A_2} \\
\\
\text{APP} \\
\frac{\Gamma \vdash M_1 : A_2 \rightarrow A \quad \Gamma \vdash M_2 : A_2}{\Gamma \vdash \text{ap}(M_1, M_2) : A}
\end{array}$$

Figure 1: Typed λ -Calculus Statics

2 Simple Types

The syntax of the language considered here is given by the following grammar:

$$\begin{aligned}
A &::= \mathbf{1} \mid \mathbf{2} \mid A_1 \times A_2 \mid A_1 \rightarrow A_2 \\
M &::= x \mid \text{yes} \mid \text{no} \mid \star \mid \langle M_1, M_2 \rangle \mid M \cdot \mathbf{1} \mid M \cdot \mathbf{2} \mid \lambda_{A_1}(x . M) \mid \text{ap}(M_1, M_2)
\end{aligned}$$

The statics is entirely standard, defining the typing judgment $\Gamma \vdash M : A$, in such a way that the structural properties are admissible. Contraction and exchange are accounted for by treating the typing context Γ as a finite set of variable typings $x_1:A_1, \dots, x_n:A_n$ in which $x_i \neq x_j$ whenever $i \neq j$. Weakening is built-in by stating all rules with an ambient typing context Γ that goes along for the ride. See Figure 1 for the definition of typing. Substitution (transitivity), which states that if $\Gamma, x:A \vdash N : B$, and $\Gamma \vdash M : A$, then $\Gamma \vdash [M/x]N : B$, is readily proved by induction on the first premise.

The dynamics is given by a transition system $M \mapsto M'$ between closed λ -terms of some type. Any closed typed term is a valid initial state. Final states are defined along with transition in Figure 2.

Theorem 1 (Preservation). *If $M : A$ and $M \mapsto M'$, then $M' : A$.*

Proof. By induction on transition. □

3 Termination Proof

The goal is to prove termination for terms of observable type:

Theorem 2 (Termination). *If $M : \mathbf{2}$, then either $M \mapsto^* \text{yes}$ or $M \mapsto^* \text{no}$.*

That is, any complete program either accepts or rejects.

Given the statement of the theorem, practically the only move available is to proceed by induction on typing. Let us consider some cases.

YES	NO	UNIT	PAIR	LAM	LFT $M \mapsto M'$
$\overline{\text{yes final}}$	$\overline{\text{no final}}$	$\overline{\star \text{ final}}$	$\overline{\langle M_1, M_2 \rangle \text{ final}}$	$\overline{\lambda_{A_1}(x . M_2) \text{ final}}$	$\overline{M \cdot 1 \mapsto M' \cdot 1}$
RHT $M \mapsto M'$		LFT-PAIR		RHT-PAIR	
$\overline{M \cdot 2 \mapsto M' \cdot 2}$		$\overline{\langle M_1, M_2 \rangle \cdot 1 \mapsto M_1}$		$\overline{\langle M_1, M_2 \rangle \cdot 2 \mapsto M_2}$	
APP $M_1 \mapsto M'_1$			APP-LAM		
$\overline{\text{ap}(M_1, M_2) \mapsto \text{ap}(M'_1, M_2)}$			$\overline{\text{ap}(\lambda_{A_2}(x . M), M_2) \mapsto [M_2/x]M}$		

Figure 2: Typed λ -Calculus Dynamics

VAR Does not apply to closed terms.

YES Immediate, as **yes final**.

NO Immediate, as **no final**.

UNIT Does not apply, not of type **2**.

PAIR Does not apply, not of type **2**.

LFT By induction, *um*

RHT By induction, *um*

FUN Does not apply, not of type **2**.

APP By induction applied to the first premise, *um*

All cases are trivial, or completely unclear.

Well, because the subterms of a term of type **2** need not have type **2**, it seems clear that it is necessary to strengthen the theorem to say something about terms of any type.

Lemma 3. *If $M : A$, then there exists N such that N final and $M \mapsto^* N$.*

The lemma suffices for the theorem because of the definition of finality for terms of type **2**. Let us consider the proof of this lemma.

VAR Does not apply to closed terms.

YES Immediate, as **yes final**.

NO Immediate, as **no final**.

UNIT Immediate, as \star final.

PAIR Immediate, as $\langle M_1, M_2 \rangle$ final.

LFT By induction there exists N such that N final and $M \mapsto^* N$. By preservation and the definition of finality N must be of the form $\langle N_1, N_2 \rangle$. By the definition of transition

$$M \cdot \mathbf{1} \mapsto^* \langle N_1, N_2 \rangle \cdot \mathbf{1} \mapsto N_1.$$

But now what?

RHT Analogous, what to do with N_2 ?

FUN Immediate, as $\lambda_{A_1}(x . M_2)$ final.

APP By induction applied to the first premise there exists N_1 such that N_1 final and $M_1 \mapsto^* N_1$. By preservation and the definition of finality N_1 must have the form $\lambda_{A_2}(x . M)$. By the definition of transition

$$\text{ap}(M_1, M_2) \mapsto^* \text{ap}(\lambda_{A_2}(x . M), M_2) \mapsto [M_2/x]M.$$

But now what?

In the projection cases the components of the pair are general terms about which nothing is known. In the application case the value of the first argument is a λ -abstraction whose body is an open term (with free variable x) about which nothing is known. This suggests strengthening the lemma by proving a property called *hereditary termination*, which is stronger than mere termination. It should have the following characteristics in order to push through the proof of the strengthened lemma below:

1. A hereditarily terminating expression of type **1** should be terminating, and hence transition to \star .
2. A hereditarily terminating expression of type **2** should be terminating, and hence transition to either **yes** or **no**.
3. A hereditarily terminating expression of type $A_1 \times A_2$ should terminate with a pair $\langle N_1, N_2 \rangle$ such that both N_1 and N_2 are hereditarily terminating.
4. A hereditarily terminating expression of type $A_2 \rightarrow A$ should terminate with a function $\lambda_{A_2}(x . M)$ such that if M_2 is hereditarily terminating of type A_2 , then $[M_2/x]M$ should be hereditarily terminating at type A .

These conditions constitute a *definition* of the property *M is hereditarily terminating at type A* , which is defined for closed $M : A$. The first two cases are given outright; the others rely on hereditary termination at constituent types of a compound type. *Thus, hereditary termination at a type is defined by induction on the structure of the type.*¹

Lemma 4. *If $M : A$, then M is hereditarily terminating at type A .*

¹For reference the type-indexed family of predicates, $\text{HT}_A(M)$, defining hereditary termination is given in Figure 3.

$$\begin{aligned}
& \text{HT}_1(M) \text{ iff } M \mapsto^* \star \\
& \text{HT}_2(M) \text{ iff } M \mapsto^* \text{yes or } M \mapsto^* \text{no} \\
& \text{HT}_{A_1 \times A_2}(M) \text{ iff } M \mapsto^* \langle M_1, M_2 \rangle \text{ and } \text{HT}_{A_1}(M_1) \text{ and } \text{HT}_{A_2}(M_2) \\
& \text{HT}_{A_1 \rightarrow A_2}(M) \text{ iff } M \mapsto^* \lambda_{A_1}(x . M_2) \text{ and } \text{HT}_{A_1}(M_1) \text{ implies } \text{HT}_{A_2}([M_1/x]M_2) \\
& \text{HT}_\Gamma(\gamma) \text{ iff } \text{HT}_A(\gamma(x)) \text{ for all } x : A \in \Gamma
\end{aligned}$$

Figure 3: Hereditary Termination, $\text{HT}_A(M)$

The proof proceeds as before by induction on typing. The cases for the constants are immediate by the definition of hereditary termination at base type.

The problematic elimination cases use the definition of hereditary termination, along with an additional property, called *head expansion*. Before stating it, let us see how it arises. Consider the rule LFT once again. By induction on the premise of the rule, $\text{HT}_{A_1 \times A_2}(M)$. By the definition of hereditary termination $M \mapsto^* \langle M_1, M_2 \rangle$ and $\text{HT}_{A_1}(M_1)$. To show $\text{HT}_{A_1}(M \cdot 1)$, observe that

$$M \cdot 1 \mapsto^* \langle M_1, M_2 \rangle \cdot 1 \mapsto M_1.$$

To complete the proof it suffices to show that hereditary termination is closed under “reverse execution”.

Lemma 5 (Head Expansion). *If $\text{HT}_A(M)$ and $M' \mapsto M$, then $\text{HT}_A(M')$.*

Proof. Immediate, because the definition of hereditary termination is defined in terms of the evaluation behavior of terms. \square

This completes the proof for the rule LFT; rules RHT and APP are handled similarly.

What about the pair and function cases?

PAIR By induction M_1 is hereditarily terminating at A_1 and M_2 is hereditarily terminating at type A_2 ; the goal is to show that $\langle M_1, M_2 \rangle$ is hereditarily terminating at type $A_1 \times A_2$. A pair is already a value (final state), so an appeal to the inductive hypothesis suffices to finish the proof.

FUN To show that $\lambda_{A_1}(x . M_2)$ is hereditarily terminating at $A_1 \rightarrow A_2$, show that whenever M_1 is hereditarily terminating at A_1 , then $[M_1/x]M_2$ is hereditarily terminating at A_2 . But what to do?

The problem now is that in the function case there is no inductive hypothesis available to give us the necessary information about the *open* term M , which has one free variable, x , in it. The lemma must be strengthened once more to account for open terms, even though the desired property applies only to closed terms.

To state the required result, define $\gamma : \Gamma$ to mean that γ is a map assigning to each variable x declared in Γ a term of the type specified in Γ . The action $\hat{\gamma}(M)$ of a substitution γ is defined inductively on the structure of M to replace each free variable x with the term assigned to it by γ .

A substitution γ is hereditarily terminating at Γ iff whenever $\Gamma \vdash x : A$, then $\gamma(x)$ is hereditarily terminating at type A .

Define $\Gamma \gg M \in A$ to mean that if γ is hereditarily terminating at Γ , then $\hat{\gamma}(M)$ is hereditarily terminating at type A . We may then state the main lemma as follows:

Lemma 6. *If $\Gamma \vdash M : A$, then $\Gamma \gg M \in A$.*

The proof is by induction on typing derivations. The critical case is the last one in the preceding attempt, for which the strengthened statement provides precisely what is needed to push the proof through. The other cases require a bit more care in handling the application of γ to the terms in question, but there are no further obstacles to worry about.

And that is Tait's Method!

Exercise 1. *If termination is required only for closed programs of answer type, and not for higher types, then a "negative" formulation of hereditary termination is sensible:*

$$\begin{aligned} HT_{A_1 \times A_2}(M) &\text{ iff } HT_{A_1}(M \cdot 1) \text{ and } HT_{A_2}(M \cdot 2) \\ HT_{A_1 \rightarrow A_2}(M) &\text{ iff } HT_{A_1}(M_1) \text{ implies } HT_{A_2}(\text{ap}(M, M_1)) \end{aligned}$$

Re-prove the termination theorem using this revised definition of hereditary termination at product and function types.

Exercise 2. *Finite sums, given by the empty type $\mathbf{0}$ and the binary sum, $A_1 + A_2$, require a "positive" formulation of hereditary termination:*

$$\begin{aligned} HT_{\mathbf{0}}(M) &\text{ iff (never)} \\ HT_{A_1 + A_2}(M) &\text{ iff } M \mapsto^* 1 \cdot M_1 \text{ and } HT_{A_1}(M_1), \text{ or} \\ &\quad M \mapsto^* 2 \cdot M_2 \text{ and } HT_{A_2}(M_2). \end{aligned}$$

Extend the proof of termination to account for sum types based on these definitions. What would be a "negative" formulation of sum types? What goes wrong?

Exercise 3. *Extend the termination proof to account for the type \mathbf{N} of natural numbers, generated by zero and successor, and interpreted by iteration, under a lazy dynamics whereby any successor is a value, regardless of the form of the predecessor. Define hereditary termination at type \mathbf{N} as the strongest property \mathcal{P} of $M : \mathbf{N}$ such that*

1. *If $M \mapsto^* \text{zero}$, then $\mathcal{P}(N)$, and*
2. *If $M \mapsto^* \text{succ}(N)$ and $\mathcal{P}(N)$, then $\mathcal{P}(M)$.*

From this definition derive a suitable induction principle to use in the proof of termination by Tait's method.

References